Index to Volume 76 of The Bryologist[1, 2]

Abietinella abietina, 407
Acarospora badiofusca, 54; glaucocarpa, 54, 395; strigata, 396; veronensis, 396
Acid phosphatases in Polytrichum commune, Electrophoretic study of, 554
Acrobolbus ochrophyllus, 514
Acrocryphaea caripensis, 486; coffaea, 151; filiformis, 496; gardneri, 486
Acrolophozia fuegiana, 519
Acroporium acestrostegium, 340; brevicuspidatum, 176; lepinei, 176; pungens, 340, 494
Actinodontium portoricense, 489
Actinogyra muehlenbergii, 335, 336
Additional mosses from Costa Rica and Mexico, 447
Additional new and noteworthy records for Hawaiian mosses, 296
Additions to the bryophyte flora of Alaska, 442; to the bryophyte flora of Mt. McKinley National Park, Alaska, 563; to the lichen flora of Arizona. I., 545; to the moss flora of Arizona. II., 439; to the moss flora of Mexico—II., 450; to the moss flora of Texas, and a new species of Isopterygium, 440
Adelanthus lindenbergianus, 529
Adelothecium bogotense, 340, 489
Agrestia hispida, 396
Ahlesia sphaerospora, 56
AHTI, TEUVO, GEORGE W. SCOTTER, and HEINO VÄNSKÄ: Lichens of the Reindeer Preserve, Northwest Territories, Canada, 48
Air pollution, 1, 333
Alaska, Additions to the bryophyte flora of, 442; Additions to the bryophyte flora of Mt. McKinley National Park, 563
Alberta, On the first 1971 foray of the American Bryological and Lichenological Society, Species collected in, Part I. Introduction and lichens, 388; Part II. Bryophytes, 403; Distributional studies on bryophytes of, 505
Alectoria chalybeiformis, 398, 546; fremontii, 398; glabra, 56, 398; implexa, 398; lanestris, 56; minuscula, 398; nigricans, 56; nitidula, 56; ochroleuca, 56; pubescens, 56, 398; sarmentosa, 209, 398; simplicior, 56; tortuosa, 209
alectorialic acid, 372
alectoronic acid, 21, 28, 29, 549
ALMEDA, FRANK, JR., and JONATHAN P. DEY: Chemical and nomenclatural notes on the Parmelia xanthina complex, 541
Aloina, Aloinella, and Crossidium (Musci), a quantitative study of, 301
Aloina **apiculata** comb. nov., 273; bifrons, 273; brevirostris, 506; hamulus, 274; longirostris, 274; pilifera, 273; rigida, 506; var. pilifera, 273; **roseae** comb. nov., 273; stellata var. pilifera, 273; sullivaniana, 273
Aloinella **andina** sp. nov., 275, 276; apiculata, 273; cucullifera, 276; hamulus, 274
Amblystegium compactum, 447; fluviatile var. **noterophilum** comb. nov., 286; juratzkanum, 254, 255, 256, 258; noterophilum, 286; riparium fo. **fluitans** comb. nov., 286; serpens, 408, 450; varium, 557
AMERICAN BRYOLOGICAL AND LICHENOLOGICAL SOCIETY, Species collected in Alberta on the first 1971 foray of, Part I. Introduction and lichens, 388; Part II. Bryophytes, 403
Amphi-Beringian flora element, Cetraria inermis (Nyl.) Krog, a new lichen species in the, 299
Amphidium lapponicum, 564
Anacamptodon cubensis, 491; splachnoides, 563
Anacolia intertexta, 483; laevisphaera, 483
Anaptychia speciosa, 399
Anastrophyllum minutum, 571
ANDEREGG, DOYLE E., GARY J. SCHROEDER, and NANCY E. SCHROEDER: Cladonia yunnana new to North America, 199; Further additions to the lichen flora of Idaho, 207
ANDERSON, LEWIS E., HOWARD A. CRUM, and WILLIAM C. STEERE: A new list of mosses of North America north of Mexico, 85
ANDERSON, LEWIS E., and BETTY E. LEMMON: Weissia sharpii, a new species from North America, with cytological and ecological correlations, 131
Andreaea, Observations on the rhizoids of, 185; alpestris, 186; blyttii, 186; nivalis, 186; rothii, 186; rupestris, 186, 474, 561
Androgynous receptacles in Marchantia polymorpha, Occurrence of, 428

[1] Compiled by William Campbell Steere, New York Botanical Garden, Bronx, New York 10458.

[2] New scientific names are printed in **boldface** type.

ANGERMAN, CARL, and MARVIN W. MEYER: Enzymes in the moss Funaria hygrometrica: phenylalanine ammonia lyase and its control, 361
Anoectangium peckii, 141; weisioides, 479
Anomobryum auratum, 203, 205; ceramiocarpum, 481; clavicaule, 481; filiforme, 481; japonicum, 203; nitidum, 203; plicatum, 481; prostratum, 481; pycnobaseum, 481
Anomodon attenuatus, 146; minor, 140; rostratus, 140; triste, 563
Anthelia juratzkana, 405
Anthoceros punctatus, 529
Anthocerotae from Southern South America, with new taxa and notes on range extensions, the Raymond E. Hatcher collection of Hepaticae and, 528
Aongstroemia bicolor, 477; fendleri, 476; tovariensis, 477; trumpfii, 477
Aplodon (Musci: Splachnaceae), Observations on the genus, 347; A. wormskioldii, 347 ff.
Apometzgeria from southern South America, and Metzgeria litoralis sp. nov., 293; A. frontipilis, 295, 529; pubescens, 295, 529
Aptychella americana, 340, 494; proligera, 494; robusta, 298
Archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, Effect of photoperiod, temperature, and plant growth hormones on initiation of, 501
Arizona, Additions to the lichen flora of, 545; Additions to the moss flora of, 439; Two new species of Xanthoparmelia from, 214
Arthonia intertexta, 56, 75; lapidicola, 56; patellulata, 401
Arthopyrenia punctiformis, 56
Arthothelium spectabile, 43
Ascidium (Ocellularia) melanostomum, 80
Astomum exserta, 142; ludovicianum, 131, 142; muhlenbergianum, 142, 558; **occidentale** sp. nov., 286; phascoides, 287, 558
Atractylocarpus longisetus, 475, var. **laxifolius** comb. nov., 474, 476, var. longisetus, 475, var. **meridense** comb. nov., 473, 476
atranorin, 21, 27, 29, 79, 230, 232, 246, 247, 248, 250, 251, 265, 370, 374, 375, 380, 382, 383, 386, 421, 542
Atrichum androgynum var. oerstedianum, 496; angustatum, 558; selwynii, 439; undulatum, 439
Aulacomnium palustre, 255, 256, 257, 259, 407, var. congestum, 445; venezuelanum, 483
Austral Hepaticae I. Pigafettoa Mass., 511; II. Evansianthus, a new genus of Geocalycaceae, 516

Bacidia bagliettoana, 56, 393; beckhausii, 56; fusca, 56; herbarum, 56; illudens, 56; lignaria, 66; muscorum, 56; obscurata, 56, 393; pallens, 56; populorum, 393; sphaeroides, 56, 393; subfuscula, 56; subincompta, 56; verecundula, 57
Baeomyces carneus, 57; roseus, 44, 57; rufus, 44, 57
Baja California, Mexico, Two new lichens (Ramalina) from, 211
barbatic acid, 78, 81, 83
Barbella conferva, 487
Barbilophozia hatcheri, 405; lycopodioides, 405
Barbula acuta, 406; brunneola, 447; campylocarpa, 479; cancellata, 557; convoluta, 406; cruegeri, 340, 450, 479, 557; ecuadoriensis, 479; erosa, 479; fallax, 406; fendleri, 479; hamulus, 274; laevigata, 447; linguaefolia, 447; mobilis, 479; moritziana, 479; rigida var. bifrons, 273, var. pilifera, 273; rufofusca, 444; spiralis, 479; sullivaniana, 273; unguiculata, 564; vinealis, 450
Bartleya ohioense, 141
Bartramia altogracilis, 483; brevifolia, 483; fendleri, 483; flavicans, 483; lineata, 483; macrodictya, 483; mathewsii, 483; moritziana, 483; muelleri, 483; nana, 483; potosica, 483; radicalis, 483; secunda, 483
Bartramidula fendleri, 483; nana, 483
Bazzania trilobata, 571
Biatorella moriformis, 57
BIRD, C. D.: Species collected in Alberta on the first 1971 foray of the American Bryological and Lichenological Society. Part I. Introduction and lichens, 388
BIZOT, M.: Contribution a la flore bryologique de Cuba (2° article), 340
BLACKMAN, A. J., G. C. BRATT, and J. A. CASHIN: Distribution and chemistry of Heterodea muelleri (Hampe) Nyl., 410
Blastenia sinapisperma, 400
Blepharidophyllum densifolium, 531
Blepharostoma trichophyllum, 405, 571
BOWERS, FRANK D., A. J. SHARP, and HAROLD ROBINSON: Additional mosses from Costa Rica and Mexico, 447
BOWLER, PETER A., and PHILIP W. RUNDEL: Two new lichens (Ramalina) from Baja California, Mexico, 211
Brachymenium andersonii, 141; globirameum, 481; melanopyxis, 481; papillosum, 482; peraristatum, 481; sipapoense, 481; smaragdinum, 481; sordidissimum, 481; wrightii, 340, var. mnioides, 481
Brachythecium albicans, 408; calcareum, 446, 564; curtum, 506; **delicatulum** sp. nov., 287; eupopuleum, 492; fendleri, 288, 292; meridense, 492; nelsonii var. **intermedium** var. nov., 288; plumosum, 492; rutabulum, 254, 255, 256, 258, 506; salebrosum, 254,

255, 256, 258, 408; starkei, 506, var. explanatum, 506, var. curtum, 506; stereopoma, 341, 450, 492; suberythrorrhizon, 288; trochalobasis, 450

BRASSARD, GUY R.: Review of K. Karczmarz, A monograph of the genus Calliergon (Sull.) Kindb., 327; and JAMES A. HANCOCK: Buxbaumia aphylla Hedw. in southeastern Newfoundland, Canada, 187

BRATT, G. C., A. J. BLACKMAN, and J. A. CASHIN: Distribution and chemistry of Heterodea muelleri (Hampe) Nyl., 410

BREIL, DAVID A.: Two bryophytes new to Virginia, 315

Breutelia brittoniae, 447; scoparia, 483; tomentosa, 340, 483

British Columbia, New and interesting macrolichens of, 366

BRODO, IRWIN M.: The lichen genus Coccotrema in North America, 260

Bruchia brevifolia, 558; drummondii, 558; sullivantii, 557

Bryoerthrophyllum recurvirostrum, 406, 435, 447; var. **dentatum** comb. nov., 288

Bryophyte flora of Alaska, Additions to, 442

Bryophytes in urban environments, Evaluation of the pollution and drought hypotheses in relation to, 1

Bryosedgwickia pallidissima, 493

Bryum andino-roseum, 482; argenteum, 206, 407, 481, var. lanatum, 340, var. normale, 481; atropurpureum, 203, 205; billiardieri, 481; caespiticium, 407; capillare, 481; ceramiocarpum, 481; chrysoblastum, 481; congestum, 482; coronatum, 340, 482; cruegeri, 340, 482; curvifolium, 482; dimorphum, 482; erythrocarpum, 185; fendleri, 482; flexisetum, 482; globirameum, 481; klinggraeffii, 183, 184, 185; leptopelma, 482; limbatum var. cubense, 340; longipedicellatum, 482; melanopyxis, 481; micropendulum, 482; pallescens, 407; papillosum, 482; peraristatum, 481; porphyroneuron, 203, 205; prostratum, 481; pseudotriquetrum, 203, 205, 407; pycnobaseum, 481; pycnopyxis, 482; ramosum, 205; riparium, 184, 185; smaragdinum, 481; sordidissimum, 481; subleucophyllum, 482; swartzianum, 482; truncorum, 340, 481; turbinatum, 558; valenciae, 482; wagneri, 482; weberaceum, 175

Bryum klinggraeffii Schimp., In vitro production of secondary gemmae on the protonema of, 183

Buellia aethalea, 57; canescens, 38; disciformis, 57, 399; epigaea, 399; immersa, 57; insignis, 57; papillata, 57; pullata, 546; punctata, 57, 399; retrovertens, 399; scabrosa, 57; spuria, 57; triphragmioides, 399; turgescens, 546; vilis, 57; zahlbruckneri, 57, 399

Buxbaumia aphylla, 187, 188, 189, 190, 565; in southeastern Newfoundland, Canada, 187; in Alaska, 565; indusiata, 444; subcylindrica, 315

Calicium abietinum, 57; populneum, 401; salicinum, 57, 401; trabinellum, 57; viride, 57, 401

California, A new Orthotrichum from northern, 437; New records and comments on lichens of, 209

Callicostella aspera, 489; chlorina, 176; depressa, 489; galipanoana, 489; meridense, 496; microcarpa, 489; nukahivensis, 176; oblongifolia, 176; pallida, 489; scabriseta, 489

Callicostellopsis meridensis, 489, 496

Calliergon cordifolium, 327; cuspidatum, 327; giganteum, 327, 408; macounii, 327, 328; obtusifolium, 327, 328; richardsonii, 327; sarmentosum var. flagellaris, 327; stramineum, 327; turgescens, 327; wickesii, 328

Caloplaca ammiospila, 57; aurantiaca, 400; borealis, 57; caesiorufella, 57; cerina, 57, 400; cinnamomea, 57; citrina, 58; discoidalis, 58; discolor, 546; ferrugineofusca, 58; fraudans, 546; holocarpa, 57, 58, 400; jungermanniae, 58, 400; leuconeura, 58; lithophila, 58; pyracea, 58; saxicola, 58; sinapisperma, 58; spitsbergensis, 58; stillicidiorum, 58, 400; tetraspora, 58; tirolensis, 58; trachyphylla, 546; variabilis, 36; wrightii, 546

calycin, 386

Calymperes aongstroemii, 175; bartramii, 479; conchophyllum, 479; donnellii, 340, 479; erosum, 479; hawaiiense, 298; lonchophyllum, 479; nashi, 341; nicaraguense, 479; perinvolutum, 479; richardii, 341, 479; venezuelanum, 479

Calymperopsis semiliber, 296

Calypogeia fissa, 571; meylanii, 405; neesiana, 443; sphagnicola, 405, 531

Calyptothecium duplicatum, 341; moritzii, 488, 497; urvilleanum, 176

Campylium chrysophyllum, 407; hispidulum, 407; stellatum, 254, 255, 258, 259, 407, var. protensum, 445, 564

Campylopodium fendleri, 476; pusillum, 341

Campylopus aerodictyon, 476; arctocarpus, 476; atratus, 476; bolivarensis, 476; breweri, 476; brittonae, 341; caracasanus, 476; chionophilus, 477; chrismarii, 476; chrysodictyon, 476; concolor, 476; crispicoma, 476; cubensis, 341; cygneus, 341; erectus, 477; exaltatus, 476; exfimbriatus, 476; falcatulus, 476; fendleri, 476; filifolius, 341,

476; flexuosus, 341, 476; fragilis var. obscurus, 341; gracilicaulis, 341; hoffmannii, 476; introflexus, 476; jamesonii, 476; lamellinervis, 476; leucogaster, 476; luteus, 476; penicillatus, 476; percurvatus, 476; polytrichoides, 341, var. altecristatus, 341; porphyreocaulis, 476; porphyreodictyon, 341, 476; praealtus, 477; pseudofilifolius, 477; richardii, 175, 477; roraimae, 476; rosulatus, 477; saxatilis, 341; sellowianus, 476; steyermarkii, 477; subulatus, 447, 448, 477; surinamensis, 477; umbellatus, 175; venezuelanum, 484
Candelaria concolor, 396
Candelariella aurella, 58, 397, 546; deflexa, 546; lutella, 58; rosulans, 207, 546; terrigena, 207; vitellina, 58, 397; xanthostigma, 58
caperatic acid, 370
Carinafolium tatei, 478
Cashin, J. A., A. J. Blackman, and G. C. Bratt: Distribution and chemistry of Heterodea muelleri (Hampe) Nyl., 410
Catagonium politum, 488
Catharinaea oerstedianum, 496; valenciae, 496
Catillaria atropurpurea, 58; glauconigrans, 393; globulosa, 393
Catoscopium nigritum, 407
Cavernularia hultenii, 210
Cephalozia bicuspidata, 405, 571; connivens, 571; media, 405, 571; patagonica, 531; pleniceps, 405
Cephaloziella byssacea, 443, 571; dusenii, 531; **gemmata** sp. nov., 528 ff.; subdentata, 405
Ceratodon purpureus, 475, 506, var. conicus, 446; semilunaris, 475; stenocarpus, 475; venezuelensis, 475
Cetraria canadensis, 209; chlorophylla, 397; commixta, 397; crispa, 299; cucullata, 58, 306, 397; delisei, 58; ericetorum, 58, 59, 299, 397; halei, 397; hepatizon, 58, 306, 397; idahoensis, 370; **inermis** comb. nov., 299 ff.; islandica, 59, 541; juniperina, 209; laevigata, 58, 59, 74; merrillii, 397; nivalis, 59, 397; pallidula, 209, 370; pinastri, 59, 397; platyphylla, 397; richardsonii, 59, 74; sepincola, 59, 299; sibirica, 59, 74; subalpina, 299; tenuifolia, 299, var. inermis, 299; tilesii, 59
Cetraria inermis (Nyl.) Krog, a new lichen species in the Amphi-Beringian flora element, 299
Cetrelia cetrarioides, 370
Chaenotheca phaeocephala, 59
Chamberlainia stereopoma, 341
Chance, Kathleen H., and Donald L. Tibbetts: A specific method for the identification of usnic acid, 208
Cheilotheca vaginata, 475
Chemical and nomenclatural notes on the Parmelia xanthina complex, 541
chemical races, 20 ff.
Chemistry of Heterodea muelleri (Hampe) Nyl., 410
Chiloscyphus fragilis, 443; hookeri, 533, var. constantifolius, 533; integrifolius, 533; pallescens, 405, 571; pallido-virens, 533; polyanthus, 405
chloroatranorin, 374, 375
Chopra, R. N., and M. S. Rawat: In vitro production of secondary gemmae on the protonema of Bryum klinggraeffii Schimp., 183; and Sneh Sood: In vitro studies on the reproductive biology of Riccia crystallina, 278
Chorisodontium wallisii, 477
Cinclidium arcticum, 432; stygium, 407
Cladina alpestris, 61; arbuscula, 394; mitis, 61, 394; rangiferina, 61, 394
Cladonia acuminata var. norrlinii, 59; alaskana, 59, 74; alpestris, 190; alpicola, 61; amaurocraea, 59; arbuscula, 54, 59, subsp. beringiana, 74; bacillaris, 59, 394; bacilliformis, 59; bellidiflora, 60; beringiana, 59; boryi, 421; botrytes, 394; cariosa, 60, 394; cenotea, 60, 394, 546; chlorophaea, 60, 370, 387, 394; coccifera, 60, 190, 395; cornuta, 60, 395, 546; crispata, 60; cristatella, 8, 335, 336, 395; cryptochlorophaea, 60; decorticata, 60; deformis, 60, var. nigripes, 60; ecmocyna, 60, 61, 74, 395; elongata, 60; fimbriata, 60, 395; floerkeana, 39; furcata, 333, 335, 336, 395, 414; gonecha, 60, 395; gracilis var. dilatata, 60, 395, var. macroceras, 60, var. **nigripes** comb. nov., 60; granulans, 61, 74; grayi, 61; lepidota, 61; macrophylla, 61; macrophyllodes, 395; merochlorophaea, 370; mitis, 54, 61, 209; multiformis, 61, 395; norrlinii, 59; pachycladodes, 420; pacifica, 209; perforata, 420; phyllophora, 61, 207, 395; pleurota, 61, 395; pocillum, 61, 395; polycarpoides, 8; polydactyla, 207; **psoromica** sp. nov., 418 ff.; pyxidata, 61, 190, 395; rangiferina, 54, 61, 190; rufescens, 41; scabriuscula, 395; scabrosa, 41; squamosa, 61; stellaris, 54, 61, 75; subfurcata, 62; subsetacea, 420; subtenuis, 545; subulata, 62, 395; symphycarpa, 60, 62; transcendens, 199, 200, var. yunnana, 199, 200; uncialis, 62, 335, 336, 395, 420, 421; venosa, 41; verticillata, 62, 207, 395; wainii, 62; yunnana, 199, 200
Cladonia psoromica, a new lichen species from western North Carolina, 418
Cladonia yunnana new to North America, 199
Cladopodiella fluitans, 405

Claopodium nervosum, 297
Clasmatocolea georgiensis, 518, 520; humilis, 533; marginata, 520; navistipula, 533; rigens, 533; vermicularis, 533
Clasmatodon parvulus, 558
Cleistostoma elegans, 525
Clidostoma rufulum, 523, 525
Climacium americanum, 140; dendroides, 407
Coccotrema antarcticum, 266; curcurbitula, 260, 266; **maritimum** sp. nov., 260, 266; **pocillarium** comb. nov., 260, 267, 268, 269
Coccotrema in North America, The lichen genus, 260
Collema auriculatum, 392; nigrescens, 392; pulposum, 39, 41; subfurvum, 546; tenax, 392; tuniforme, 62; undulatum, 546
Cololejeunea biddlecomiae, 571
Colura naumanni, 514
connorstictic acid, 21 ff.
Conocephalum conicum, 571
Conomitrium biareolatum, 475; caripense, 475; goebelii, 475; latiusculum, 475; subulatifolium, 475; trachelymum, 475
Conostomum pentastichum, 483; tetragonum, 407
conphysodic acid, 375
conpsoromic acid, 214
constictic acid, 22 ff., 230, 232, 233, 235, 238, 242, 244, 265, 268, 385, 386
Contribución a la bryoflora de Venezuela II, 559
Contribution a la flore bryologique du Cuba (2° article), 340
Cornicularia aculeata, 62; divergens, 62; muricata, 62; umhausensis, 370
Costa Rica and Mexico, Additional mosses from, 447
Costesia spongiosa, 536
Cratoneuron falcatum var. **williamsii** comb. nov., 288; filicinum, 407, 492; papillosum, 288; williamsii, 288
CROSBY, MARSHALL R.: Recent literature on mosses, 316, 456, 578
Crossidium aberrans, 277, fo. **epilosum** fo. nov., 289; peruvianum, 273; rosae, 273; spatulaefolium, 277
Crossidium (Musci), A quantitative study of, 301
Crossomitrium goebelii, 489; patrisiae, 489; phragmidiaceum, 489; tenellum, 489
CRUM, HOWARD A.: Review of Jane Taylor and Alan P. Underhill, Liverworts. A photographic study, 590; and WILLIAM C. STEERE and LEWIS E. ANDERSON: A new list of mosses of North America north of Mexico, 85
Cryphaea caripensis, 486; filiformis, 144 ff., 486, 496; floridensis, 144 ff.; glomerata, 144 ff., 557, var. scabra, 144 ff.; langloisii, 152; nervosa, 144 ff.; patens, 486; pendula, 144 ff.; ravenelii, 144 ff.; reticulata, 486; rufescens, 523, 525
Cryphaea in North America north of Mexico, A revision of the genus, 144
Cryptopodium hookeri, 483; jamesonii, 483
Ctenidiadelphus cylindricarpus, 495
Ctenidium malacodes, 495
Cuba, Contribution a la flore bryologique de, 340
CULBERSON, CHICITA F., and MASON E. HALE, JR.: 4-O-demethylnotatic acid, a new depsidone in some lichens producing hypoprotocetraric acid, 77
CULBERSON, WILLIAM LOUIS: The Parmelia perforata group: Niche characteristics of chemical races, speciation by parallel evolution, and a new taxonomy, 20; Recent literature on lichens, 217, 321, 461, 584
Cupressina sanguiseta, 494
Cyclodictyon albicans, 341, 489; albicaule, 341; amnigenum, 489; capillatum, 489; cuspidatum, 341, 489; fendleri, 489; jagianum, 489; lindigianum, 341; nivale, 489; varians, 489
Cynodontium schistii, 564
Cyphelium inquinans, 62, 401; tigillare, 62, 401
Cytological and ecological correlations, Weissia sharpii, a new species from North America, 131
Cytological studies on some West Himalayan mosses—III, 202

Dactylina arctica, 62; ramulosa, 62
Daltonia fendleri, 489; filiformis, 149; glomerata, 155; heteromalla, 144, 152, 155; longifolia, 489; nervosa, 156, 159; stenophylla, 489; tenuifolia, 490
DELGADILLO M., CLAUDIO: A new species, nomenclatural changes, and generic limits in Aloina, Aloinella and Crossidium, 271; A quantitative study of Aloina, Aloinella, and Crossidium, 301; and H. ROBINSON: Neosharpiella, a new genus of Musci from high elevations in Mexico and South America, 536
Dendriscocaulon umhausense, 370
Dendropogon rufescens, 522, 523
Dendropogonella Britt., A review of the monotypic genus, 521
Dendropogonella rufescens, 521 ff.
Dermatocarpon intestiniforme, 62; lachneum, 62; miniatum, 42, 401; plumbeum, 546; rufescens, 62
Desmatodon garberi, 341; latifolius, 406; obtusifolius, 292, 427; sprengelii, 341; stomatodontus, 341

Desmaziera sinensis, 70
DEY, JONATHAN P.: Cladonia psoromica, a new lichen species from western North Carolina, 418; Two lichens (Parmeliae) new to the Southern Appalachian Mountains, 548; and FRANK ALMEDA, JR.: Chemical and nomenclatural notes on the Parmelia xanthina complex, 511
Dichelyma falcatum, 445
Dicnemoloma pallidum (Hook.) Wijk & Marg., Unusual sporocytes in, 178
Dicnemoloma pallidum, 178 ff.
Dicranella bicolor, 477; harrisii, 341; heteromalla, 557; hilariana, 477; perottetii, 477; rigidula, 444; rufiseta, 175; subinclinata, 477; subulata, 406; tovariensis, 477; trumpffii, 477
Dicranodontium denudatum, 296, 297, 477; pulchroalare, 477
Dicranoloma rufifolium, 175
Dicranoweisia crispula, 406
Dicranum aerodictyon, 476; asperrimum, 477; chionophilum, 477; crispicoma, 476; ecaudatum, 477; erectum, 477; exaltatum, 476; densum, 477; fendleri, 476; flagellare, 406, 450; fragilifolium, 406; frigidum, 477; fuscescens, 190, 406; herminieri, 477; longisetum, 476, var. laxifolium, 476; montanum, 444; polysetum, 406, 561; porphyreocaule, 476; porphyreodictyon, 476; proliferum, 476; pseudofilifolium, 477; scoparium, 140, 406, 477; spurium, 190; undulatum, 406, 564; zygodonticarpum, 476
Didymodon alpigenus, 288; angustifolius, 479; diaphanobasis, 480; fuscoviridis, 447, 448; godmanianus, 447, 448; inundatus, 479; johansenii, 446; juniperinus, 447, 448; moritzianus, 479; recurvirostris var. dentatus, 288; rigidulus, 435, 447, 448; rubellus var. dentatus, 288
diffractaic acid, 410, 412, 413
Diphyscium cumberlandianum, 141; fendleri, 496
Diplophyllum apiculatum, 571; obtusifolium, 405
Diploschistes bryophilus, 62, 546
Dirina repanda, 34, 37
Distichium capillaceum, 289, 406, var. **curvatum** var. nov., 289; inclinatum, 289, 406
Distichophyllum cuspidatum, 176
Distribution and chemistry of Heterodea muelleri (Hampe) Nyl., 410
Distributional records of rare or infrequently collected species of mosses in Pennsylvania, 560
Distributional studies on bryophytes of Alberta, 505
Ditrichum flexicaule, 406, var. densum, 444, var. sterile, 564; giganteum, 444, 564; pallidum, 557, 558; rufescens, 475; steyermarkii, 447, 448; tortile, 178
divaricatic acid, 306, 308, 309
DNA, 361 ff.
Drepanocladus aduncus, 408, var. kneiffii, 439, var. polycarpus, 254 ff., 408; exannulatus, 408, 440; revolvens, 408; sendtneri, 408; uncinatus, 408
Drepanophyllum duidense, 482; fulvum, 482; viride, 482
Drought hypotheses in relation to lichens and bryophytes in urban environments, 1
Drummondia pursellii Vitt & Crum, A second report for, 560

Ectropothecium aeruginosum, 495; apiculatum, 495; cupressoides, 495; globitheca, 495; rutilans, 494; sodale, 176
Editorial, 216, 573
Effect of photoperiod, temperature, and plant growth hormones on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
Electrophoretic study of acid phosphatases in Polytrichum commune, 554
Encalypta procera, 561; rhaptocarpa, 406; streptocarpa, 561; vulgaris, 406, 564, fo. pilifera, 444
Endocarpon pusillum, 62
endocrocin, 80, 370
ENGEL, JOHN J.: The Raymond E. Hatcher collection of Hepaticae and Anthocerotae from southern South America, with new taxa and notes on range extensions, 528; and YUKINOBU KUWAHARA: Metzgeria litoralis sp. nov. and Apometzgeria from southern South America, 293; and R. M. SCHUSTER: Austral Hepaticae I. Pigafettoa Mass., 511; II. Evansianthus, a new genus of Geocalycaceae, 516
Entodon aurescens, 493; columnaris, 493; complicatus, 493; distans, 493; hampeanus, 493; jamesonii, 493; macropodus, 493; myurus, 493; pallescens, 493; pallidissimus, 493; seductrix, 140
Entosthodon acutifolius, 481; spathulifolius, 536; trumpffii, 481
Enzymes in the moss Funaria hygrometrica: phenylalanine ammonia lyase and its control, 361
Ephebe lanata, 62
Epiphyllous Hepaticae in Louisiana, 201
Epipterygium orbifolium, 482; wrightii, 496
Erpodium acrifolium, 440; coronatum, 484; cubense, 341; domingense, 341; latifolium, 484

Erythrodontium brasiliense, 497; cylindricaule, 493; julaceum, 497; longisetum, 493; moritzii, 497; squarrosum, 493
ESSLINGER, THEODORE L.: Nomenclatural notes on some members of Parmelia section Melanoparmelia, 306; Physcia luganensis new to North America, 421
Eucamptodontopsis pilifera, 477; tortuosa, 477
Eurhynchium praelongum, 450; pulchellum, 255, 256, 408; regnellii, 492
Eustichia spruceana, 482
Evansianthus, A new genus of Geocalycaceae, 516
Evansianthus georgiensis gen. et comb. nov., 516 ff.
Evernia divaricata, 398; mesomorpha, 62, 74, 398; perfragilis, 62, 74
evernic acid, 83
Exodictyon nadaudii, 175

Fabronia glauca, 491; perimbricata, 491; polycarpa, 491; pusilla, 435; wrightii, 341
far-red light, 361
Fissidens amoenus, 474; angustifolius, 474; aoraiensis, 175; aporrocheilos, 474; asplenioides, 341, 473, 474; biareolatus, 475; bishopii, 297; brachypus, 475; bushii, 561; caripensis, 475; clarkii, 175; cristatus, 558; diplodus, 341, 447, 448, 450, 451, 475; dissitifolius, 341; **duryae** sp. nov., 340 ff.; elegans, 475; elimbatus, 203, 206; exilis, 561; fasciculatus, 343; fendleri, 475; garberi, 341; genunervis, 473, 474; goebelii, 475; hyalinus, 561; imbricatus, 343; inaequalis, 475; inclinis, 475; intermedius, 475; kegelianus, 343, 475; kilaueae, 297; latiusculus, 475; limbatus, 292; mollis, 475; monostictus, 343; moritzianus, 475; muriculatus, 341, 451, 475; obtusifolius var. kansanus, 290, var. **marginatus** var. nov., 289; oahuensis, 297; ocellatus, 343; pennula, 475; platybryoides, 343; plumosus, 343; polypodioides, 341, 475; prionodes, 475; radicans, 341, 475; reticulosus, 475; scariosus, 475; secundatus, 497; secundulus, 475; semimarginatus, 475; similiretis, 341; steerei, 475; sublimbatus, 451; subulatifolius, 475; sylvaticus, 202, 205; taxifolius, 296, 297, 561; trachelyma, 475; validicostatus, 475; vardei, 341, 447, 448; ventricosus, 208; veracruzensis, 447, 448; viridulus, 558; weirii, 341, 447, 448, var. insertus, 447, 448; yucatanensis, 343, 475
FLORSCHÜTZ, P. A.: Review of Geneva Sayre, Cryptogamae exsiccatae, Part IV. Bryophyta, 326
†FLOWERS, SEVILLE: Mosses of Utah new to science, 286
Fontinalis filiformis, 151
Forsstroemia ravenelii, 159; trichomitria, 146, 562
Fossombronia foveolata, 443, 564
Frullania asagrayana, 571; bolanderi, 571; boveana, 533; brittoniae, 572; eboracensis, 572; inflata, 505, 506; lobulata, 533; magellanica, 533; patagonica, 533
FULFORD, MARGARET: Recent literature on hepatics, 452, 574
Fulgensia bracteata, 400, 546; fulgens, 400
fumarprotocetraric acid, 370
Funaria acutifolia, 481; americana, 557; calvescens, 343, 481; flavicans, 557; hygrometrica, 356 ff., 361 ff., 406, 423 ff., 451, 481, var. calvescens, 343, 481; macrospora, 481; trumpffii, 481
Funaria hygrometrica, A time-course of sporophyte development in, 356; Phenylalanine ammonia lyase and its control by enzymes, 361; On the functioning of stomates in, 423
Functioning of stomates in Funaria, On the, 423
Further additions to the lichen flora of Idaho, 207

Gackstroemia magellanica, 533; schwabei, 533
galbinic acid, 21, 29
GARNER, DOROTHY, and DOMINICK J. PAOLILLO, JR.: A time-course of sporophyte development in Funaria hygrometrica Hedw., 356; On the functioning of stomates in Funaria, 423
Garovaglia powellii, 176
Geocalyx graveolans, 405, 572
Germinability of immature spores in Polytrichum, 163
glomellic acid, 308, 309
glomelliferic acid, 308, 309
Gloniopsis levantica, 31
Glossadelphus abortivapicus and Rhaphidostichum pustulatum, new species of the Sematophyllaceae from Hawaii, 310
Glossadelphus **abortivapicus** sp. nov., 310 ff.; andersonii, 141; baldwinii, 313; chrysobasilaris, 298; limnobioides, 297; zollingeri, 297
Glypholecia scabra, 63
granite outcrop pioneer ecosystems, Nitrogen fixation on, 196
Graphis elegans, 209; scripta, 44; striatula, 209
Grimmia affinis, 444; anodon, 406, 444; apocarpa var. ambigua, 444, 564, var. stricta, 140; calyptrata, 406; laevigata, 185, 187, 196, 197, 198; mollis, 444; ovalis, 480
Groutiella apiculata, 343; husnotii, 484; mucronifolia, 344, 484; tomentosa, 484; undosa, 344, 497; wagneriana, 344, 484
GUERKE, WAYNE R.: Epiphyllous Hepaticae in Louisiana, 201

GUNDERSON, GERALD O.: Hepatics from southwestern Wisconsin, 571
Gyalecta jenensis, 63; kukriensis, 63
Gymnostomum calcareum, 480; recurvirostrum, 480, 564; venezuelense, 480
gyrophoric acid, 230, 232, 245, 246, 248, 250, 251, 306, 307, 308, 309, 380, 382, 383, 541, 542, 551, 553, 554
Gyroweisia papillosa, 447, 448

Haematomma lapponicum, 207
Hageniella pacifica, 297
HALE, MASON E., JR., and CHICITA F. CULBERSON: 4-O-demethylnotatic acid, a new depsidone in some lichens producing hypoprotocetraric acid, 77
HAMPTON, RAYMOND E.: Photosynthetic pigments in Peltigera canina (L.) Willd. from sun and shade habitats, 543
HANCOCK, JAMES A., and GUY R. BRASSARD: Buxbaumia aphylla Hedw. in southeastern Newfoundland, Canada, 187
Haplocladium microphyllum, 296, 297, 344, 491
Haplodon wormskioldii, 347 ff.
Haplohymenium triste, 297, 563
Harpanthus scutatus, 572
HATCHER, RAYMOND E., collection of Hepaticae and Anthocerotae from southern South America, with new taxa and notes on range extensions, 528
Hawaiian mosses, Additional new and noteworthy records for, 296
Hawaiian species of the Pottiaceae, Notes on some, 192
HÉBANT, CHARLES, and GUY KAHLEM: Electrophoretic study of acid phosphatases in Polytrichum commune, 554
Hedwigia albicans, 485; ciliata, 485, 506, 557; nivalis, 485
Hedwigidium imberbe, 485
Helicophyllum torquatum, 344
Helodium blandowii, 255, 256, 257, 258, 259, 407; paludosum, 563
Hepaticae and Anthocerotae from southern South America, with new taxa and notes on range extensions, the Raymond E. Hatcher collection of, 528
Hepatics from southwestern Wisconsin, 571
Hepatics, Recent literature on, 452, 574
HERMANN, FREDERICK J.: Additions to the bryophyte flora of Alaska, 442; Additions to the bryophyte flora of Mt. McKinley National Park, Alaska, 563; and ELVA LAWTON: A new Orthotrichum from northern California, 437
Herzogobryum erosum, 533; teres, 519
Heterodea muelleri (Hampe) Nyl., Distribution and chemistry of, 410
Heterodea muelleri, 410 ff.
HOE, W. J.: Additional new and noteworthy records for Hawaiian mosses, 296; Glossadelphus abortivapicus and Rhaphidostichum pustulatum, new species of the Sematophyllaceae from Hawaii, 310; Notes on some Hawaiian species of the Pottiaceae, 192
Holomitriopsis laevifolia, 478
Holomitrium arboreum, 477; crispulum, 477; cylindraceum, 175; lutescens, 477; marginatum, 344; moritzianum, 477; pulchellum, 447, 448
Homalia membranacea, 490
Homaliodendron flabellatum, 312
Hookeria acutifolia, 344; amnigena, 489; aspera, 489; bipinnata, 489; cuspidata, 489; divaricata, 490; fendleri, 489; galipanoana, 489; goebelii, 490; jagiana, 489; membranacea, 490; meridensis, 489; muelleri, 490; nivalis, 489; pallens, 489; pallescens, 490; pernutans, 490; philonotula, 490; pilotrichelloides, 491; plumicaulis, 490; radiculosa, 493; roraimae, 490; scabriseta, 489; subenervis, 490
Hookeriopsis crispa, 490; cruegeriana, 490; falcata, 344, 490; guadalupensis, 344; incurva, 344, 490; maguirei, 490; parkeriana, 490; pernutans, 490; plumicaulis, 490; ptari-tepuiensis, 490; steyermarkii, 490; subfalcata, 490; undata, 490; venezuelensis, 490
Hormones, Effect of plant growth, on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
Hyalolepidozia bicuspidata, 533
Hydrogen ion concentrations, The effect of, in simulated rain on the moss Tortula ruralis (Hedw.) Sm., 168
Hydropogon fontinaloides, 485
Hydropogonella gymnostoma, 485
Hygroamblystegium meridense, 492; noterophilum, 286
Hygrohypnum bestii, 446; luridum fo. **serratum** fo. nov., 290, fo. **tenue** fo. nov., 290; pelichucense, 447, 448; tequendamense var. subfalcatum, 492
Hygrolembidium isophyllum, 519, 533
Hylocomium pyrenaicum, 408, 506; splendens, 255 ff., 408
Hymenostomum caripensis, 480; involutifolium, 480; microstomum, 142; tortile, 135, 137
Hymenostylium recurvirostre, 344, 451
Hyophila arborea, 480; involuta, 480; involutifolia, 480; melanostoma, 480; microcarpa, 480; tortula, 480
Hypnella cymbifolia, 490; leptorrhyncha, 490;

pallescens, 490; philonotula, 490; pilifera, 490; pilotrichelloides, 491
Hypnodendron vescoanum, 175
Hypnum aemulans, 74; aeruginosum, 495; amabile, 495; campylocarpum, 492; chrysostegum, 494; concavum, 495; cupressiforme, 7, 408, var. subjulaceum, 446, 564; delicatulum, 492; diminutivum, 495; elegantulum, 495; eupopuleum, 492; eutrypherum, 493; frontinoae, 491; galipense, 495; hamata, 495; lindbergii, 408; meridense, 492; mirabile, 495; muriculatum, 491; nanopolymorphum, 495; nigricans, 487, var. pendula, 488; noterophilum, 286; obliquefolium, 494; pauperum, 491; polypterum, 344; pratense, 408; pseudoprotensum, 492; revolutum, 408; riparium, 286; rutilans, 494; schlimii, 494; schraderi, 493; subperspicuum, 493; subscabrum, 494; tovariense, 494; trachynotum, 493; trichostegum, 494; vaucheri, 408
Hypodontium pomiforme, 497
Hypogymnia atrofusca, 371 ff.; austerodes, 63, 74, 371 ff.; bitteri, 63, 74, 371 ff., 397, 546; duplicata, 371 ff.; encausta, 372; enteromorpha, 371 ff., 397; imshaugii, 371 ff., var. inactiva, 374; **inactiva** comb. nov., 366, 371 ff.; intestiniformis, 207, 372, var. atrofusca, 372; **krogii** sp. nov., 366, 371 ff.; metaphysodes, 372 ff.; physodes, 7, 8, 9, 13, 63, 371 ff., 397; pseudophysodes, 371 ff.; subobscura, 371 ff., 379; tubulosa, 210, 371 ff., 397; vittata, 371 ff.
hypoprotocetraric acid, 77 ff., 214
Hypopterygium flavescens, 491; laricinum, 491; nivale, 491; rigidulum, 491; sylvaticum, 491; tahitense, 176; tamariscinum, 344, 491, 559

Icmadophila ericetorum, 44, 63, 396
Idaho, Further additions to the lichen flora of, 207
Identification of usnic acid, A specific method for, 208
Initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
In vitro production of secondary gemmae on the protonema of Bryum klinggraeffii Schimp., 183
In vitro studies on the reproductive biology of Riccia crystallina, 278
Isodrepanium lentulum, 344, 490
Isopaches hellerianus, 405
Isopterygium, Additions to the moss flora of Texas, and a new species of, 440
Isopterygium alstonii, 493; eutrypherum, 493; **homomallifolium** sp. nov., 440 ff.; irelandii, 493; micans, 344; minutirameum var. brevifolia, 176; pulchellum, 440; rufisetulum, 344; tenerum, 344, 493
Isotachis grossidens, 533; humectata, 533

Jaegerina scariosa, 486
JAHNS, H. M.: The trichogynes of Pilophorus strumaticus, 414
Jamesoniella autumnalis, 572; colorata, 533
JOHNSEN, ARDITH B.: Additions to the moss flora of Arizona II, 439
JORDAN, WILLIAM PAUL: The genus Lobaria in North America north of Mexico, 225
Jungermannia lanceolata, 572

KAHLEM, GUY, and CHARLES HÉBANT: Electrophoretic study of acid phosphatases in Polytrichum commune, 554
KASS, LEE B., and DOMINICK J. PAOLILLO, JR.: The germinability of immature spores in Polytrichum, 163
KERSHAW, KENNETH A.: Review of J. Levitt, Responses of plants to environmental stresses, 328
KOEVENIG, JAMES L.: Effect of photoperiod, temperature, and plant growth hormones on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
KRAL, ROBERT: Review of Arthur Cronquist, Arthur H. Holmgren, Noel H. Holmgren, and James L. Reveal, Intermountain flora, 502
KROG, HILDUR: Cetraria inermis (Nyl.) Krog, a new lichen species in the Amphi-Beringian flora element, 299; On Umbilicaria pertusa Rass. and some related lichen species, 550
KUMAR, S. S.: Cytological studies on some West Himalayan mosses—III, 202
KUWAHARA, YUKINOBU: Further notes on the production of vegetative thallus structures by female involucres of Metzgeria, and a new species of Metzgeria, 566; and JOHN J. ENGEL: Metzgeria littoralis sp. nov. and Apometzgeria in southern South America, 293

Lasallia chiriquiensis, 551; papulosa, 553; pennsylvanica var. pertusa, 551; pertusa, 551, 553; pustulata, 551, 553, var. papulosa, 553
Laurera purpurina, 40
LAWTON, ELVA, and F. J. HERMANN: A new Orthotrichum from northern California, 437
LEBLANC, FABIUS, and DHRUVA N. RAO: Evaluation of the pollution and drought hypotheses in relation to lichens and bryophytes in urban environments, 1

Lecanora alboradiata, 63; allophana, 396; apochraeoides, 63; badia, 63, 396, 546; basaltica, 63; cadubriae, 396; castaneoides, 63; chlarotera, 38; cinerea, 63; coilocarpa, 63, 396; conizaeoides, 7, 8, 12, 13, 16; crenulata, 63; discoensis, 63; dispersa, 9, 12, 13, 63; disserpens, 63; epibryon, 63, 396; expallens, 64; fuscescens, 65; garovaglii, 546; hageni, 64, 396; impudens, 396; kariana, 64; melanaspis, 64, 546; melanophthalma, 64, 546; pachythallina, 64; palanderi, 64, 74; permutata, 64; polytopa, 64; rolleana, 396; rugosella, 396, 546; rupicola, 396; rykaipiae, 64; saligna, 64; sambuci, 64; subintricata, 64; symmicta, 64; torrida, 64; umbrina, 64; urceolaria, 67; varia, 396; verrucosa, 67, 207; wisconsinensis, 64
lecanoric acid, 307, 541
Lecidea aeruginosa, 64; albohyalina, 56, 65; atrata, 65; atrobrunnea, 393; botryosa, 65; carpathica, 547; confluens, 65; coarctata, 41; crassipes, 65; crustulata, 65; decipiens, 65, 393; deustata, 65; dicksonii, 65, 393; dispersa, 13; elaeochroma, 35, var. soralifera, 66; epiiodiza, 65; erythrophaea, 65; euphorea, 66; flexuosa, 64; friesii, 393; fuscescens, 65, 393; glomerulosa, 393; granulosa, 65, 393; icterica, 547; jurana, 65; lapicida, 65, 393; limosa, 65; lithophila, 65; macrocarpa, 65, 394; magnussoniana, 65; marginata, 65; phaeopelidna, 65; ramulicola, 65; rubiformis, 65, 394; sanguinaria var. melinodes, 66; scalaris, 65, 394; somphotera, 65; sorediata, 64; sphaerella, 65; stigmatea, 394; subcandida, 65; symmicta, 64, 394; templetonii, 65; tornoensis, 66; turgidula, 66; uliginosa, 66; vernalis, 66, 394; vorticosa, 66
Lecidella elaeochroma, 66; enteroleuca, 66; flavisorediata, 66; glomerulosa, 66; pulveracea, 66; stigmatea, 66; wulfenii, 66
Leiocolea gillmanii, 405; rutheana, 405
Leiomela bartramioides, 483; lineata, 483; subbrevifolia, 483
LEMMON, BETTY E., and LEWIS E. ANDERSON: Weissia sharpii, a new species from North America, with cytological and ecological correlations, 131
Lepidolaena menziesii, 533
Lepidopilidium divaricatum, 490; purpurissatum, 490, 497
Lepidopilum aureofulvum, 344, 490; biductulosum, 490; diaphanum, 490; divaricatum, 497; goniothecium, 490; haplociliatum, 447, 448, 490; leiomitrium, 490; membranaceum, 490; mnioides, 490; muelleri, 490; polytrichoides, 490; purpurissatum, 490, 497; scabrisetum, 490; steyermarkii, 491; subpolytrichoides, 491
Lepidozia chordulifera, 533; laevifolia, 533; reptans, 405, 572
Lepraria chlorina, 547; incana, 7, 12; membranacea, 66, 401; neglecta, 66, 207, 401
Leptobryum pyriforme, 254 ff., 407
Leptodictyum riparium, 492, var. fluitans, 286
Leptodon australis, 434; smithii, 434 ff.; trichomitrion, 562
Leptodon smithii (Musci, Neckeraceae), a genus new to North America, 434
Leptodontium cirrhifolium, 480; filicola, 480; filescens, 480; flexifolium, 451, 480; longicaule var. microruncinatum, 480; pungens, 480; subcirrhifolium, 480; ulocalyx, 480; viticulosoides var. panamense, 480
Leptogium dendriscum, 383; furfuraceum, 392; lichenoides, 66, 392; saturninum, 66, 392; sinuatum, 66; tenuissimum, 66
Leptohymenium julaceum, 493
Leptophyllopsis **irregularis** comb. nov., 528, 533; laxus, 533
Leptoscyphus abditus, 516, 520; expansus, 533
Leptotheca costaricensis, 483
Leptotrema arecae, 79
Leptotrichum plumosum, 475
Lepyrodon tomentosus, 486
Lepyrodontopsis trichophylla, 344, 492
Lescuraea radicosa, 407
Leskea angustata, 491; nigricans, 487; palisotii, 487; remotifolia, 487
Leskeella **arizonae** comb. nov., 290, fo. **flagellifera** fo. nov., 290; cyrtophylla, 291; tectorum, 291, 447, 448, fo. **cyrtophylla** stat. nov., 291, subsp. cyrtophylla, 291, fo. **flagellifera** stat. nov., 291
Leskeodon longipilus, 447, 448; parvifolius, 401
Letharia columbiana, 398; vulpina, 398
LETROUIT-GALINOU, MARIE-AGNÈS: Les asques des lichens et le type archaeascé, 30
Leucobryum albicans, 344; albidum, 344, 478, 557; antillarum, 478; brevifolium, 478; crispum, 344, 478; flavomucronatum, 478; giganteum, 344, 478; laevifolium, 478; longifolium, 478; martianum, 344, 478, 559; megalophyllum, 478; microcarpum, 478; polakowskyi, 344, 478; vulgare, 497
Leucodon brachypus var. andrewsianus, 562; domingensis, 486; julaceus, 557, 558, 562; pacificus, 175; sciuroides, 562; suborthostichus, 486
Leucodontopsis geniculata, 486
Leucoloma albulum, 344; asperrimum, 477; cruegerianum, 477; ecaudatum, 477; mariei, 477; mollissimum, 477; procumbens, 477; schwaneckeanum, 344; serrulatum, 344, 477; tortellum, 478

Leucomium aneurodictyon, 176; attenuatum, 491; compressum, 491; lignicola, 491
Leucophanes glaucum, 296, 297; nukahivense, 175
Lichen crocatus, 385; duplicatus, 374; physodes, 378; retiger, 379
Lichen flora of Idaho, Further additions to, 207
Lichen genus Coccotrema in North America, 260
Lichen products and chemical substances: alectorialic acid, 372; alectoronic acid, 21, 28, 29, 549; atranorin, 21, 27, 29, 79, 230, 232, 246, 247, 248, 250, 251, 265, 370, 374, 375, 380, 382, 383, 386, 421, 542; barbatic acid, 78, 81, 83; calycin, 386; caperatic acid, 370; chloroatranorin, 374, 375; connorstictic acid, 21 ff.; conphysodic acid, 375; conpsoromic acid, 214; constictic acid, 20 ff., 230, 232, 233, 235, 238, 242, 244, 265, 268, 385, 386; diffractaic acid, 410, 412, 413; divaricatic acid, 306, 308, 309; endocrocin, 80, 370; evernic acid, 83; fumarprotocetraric acid, 370; galbinic acid, 21, 29; glomellic acid, 308, 309; glomelliferic acid, 308, 309; gyrophoric acid, 230, 232, 245, 246, 247, 248, 250, 251, 306, 307, 308, 309, 380, 382, 383, 541, 542, 551, 553, 554; hypoprotocetraric acid, 77 ff., 214; lecanoric acid, 307, 541; merochlorophaeic acid, 370; methyl hypoprotocetrarate, 79; methyl notatate, 79; 4-O-demethylbarbatic acid, 78, 81, 83; 4-O-demethylnotatic acid, 77, 78, 79, 80, 81, 83; methyl 4-O-demethylnotate, 79; 4-O-methylgyrophoric acid, 230, 245, 247, 248, 250, 251, 382; methyl 2',4-di-O-methylhypoprotocetrarate, 79; methyl 4-O-methylhypoprotocetrarate, 79; 4-O-methylhypoprotocetraric acid, 77, 78, 79, 80, 81, 83; methyl 2',-O-methylnotate, 79; norlobaridone, 307; norobtusatic acid, 77, 78, 81, 83; norstictic acid, 20 ff., 79, 230, 232, 235, 238, 242, 244, 265, 306, 385, 386; notatic acid, 77, 78, 79, 80, 83; obtusatic acid, 77, 78, 81, 83; olivetoric acid, 380, 541; orsellinic acid, 83; parietin, 386; perlatolic acid, 308, 309, 370; physodalic acid, 371, 372, 374, 375; physodic acid, 371, 372, 374, 375; protocetraric acid, 375, 380, 549; protolichesterinic acid, 299; protolichesterinic acid, 541, 542; psoromic acid, 214, 420, 421; pulvinic acid, 385, 386; pulvinic dilactone, 385, 386; rhizonic acid, 83; salazinic acid, 21, 29, 211, 380, 382, 549; scabrosin, 383; scrobiculin, 230, 232, 238, 251; squamatic acid, 200, 420, 421; stenosporic acid, 306, 308; stictic acid, 20 ff., 230, 232, 233, 235, 238, 244, 265, 268, 383, 385, 386; tenuorin, 230, 232, 233, 241, 245, 383, 386; thelephoric acid, 230, 235, 236, 237, 379; usnic acid, 80, 81, 208 ff., 213, 214, 232, 233, 238, 242, 370, 420, 542; zeorin, 383
Lichens and bryophytes in urban environments, 1
Lichens (Parmeliae) new to the Southern Appalachian Mountains, 548
Lichens of California, New records and comments on, 209
Lichens (Ramalina) from Baja California, Mexico, 211
Lichens, Recent literature on, 217, 321, 464, 584
Lichens, Sensitivity of to sulfur dioxide, 333
Lindigia trichomitria, 487, 491
linitins, 230, 233, 241
literature on hepatics, Recent, 452, 574
literature on lichens, Recent, 217, 321, 464, 584
literature on mosses, Recent, 316, 456, 578
Lobaria in North America north of Mexico, 225
Lobaria amplissima, 251; hallii, 226 ff.; kurokawae, 227 ff.; laetevirens, 34, 251; linita, 227 ff., 393, var. ochroleuca, 241; lobulifera, 244; oregana, 227 ff.; pseudopulmonaria, 227 ff.; pulmonaria, 226 ff.; quercizans, 226 ff.; ravenelii, 226 ff.; retigera, 227 ff., 379; scrobiculata, 226 ff.; tenuis, 227 ff.
Lopadium pezizoideum, 66
Lophocolea austrigena, 533; georgiensis, 518; heterophylla, 504, 572; lenta, 533; leptantha, 533; minor, 405, 572; muricata, 534; navistipula, 519; striatella, 534; sylvatica, 534; textilis, 534
Lophozia alpestris, 572; barbata, 572; badensis, 427; excisa, 534; hatcheri, 534; incisa, 405, 572; longidens, 405; obtusa, 405; porphyroleuca, 405; ventricosa, 405, 572, var. silvicola, 564
Louisiana, Epiphyllous Hepaticae in, 201

Macrocoma hymenostomum, 484
Macrolichens of British Columbia, New and interesting, 366
Macromitrium acutissimum, 484; apiculata, 484; cirrosum, 484, var. cirrosum, 344, var. jamaicense, 344; contextum, 484; densifolium, 497; dussii, 484; eurymitrium, 175; fendleri, 484; fragilis, 484; galipense, 484; guatemalense, 484; harrisii, 344; laevifolium, 484; leprieurii, 484; longifolium, 484, var. brevifolium, 484; macrothele, 484; paucidens, 484; pellucidum, 484; perichaetiale, 484; podocarpi, 484; proliferum, 484; punctatum, 484; pyriforme,

484; retusulum, 484; rhaphidophyllum, 484; richardii, 484; runcinatella, 484; schwaneckeanum, 344; serrulatum, 485; stellatum, 484, 497; stellulatum, 484; stolonigerum, 484; stricticuspis, 484; subnitidum, 485; subpaucidens, 485; swainsonii, 485; ulophyllum, 485; venezuelense, 485; wagnerianum, 484
Macrothamnium macrocarpum, 298
MAGILL, ROBERT E.: Notes on Texas mosses II, 557
Manitoba, Studies on the bryophytes of southern. VII, 252
MANUEL, MONTE G.: Additions to the moss flora of Mexico—II, 450; A second report for Drummondia pursellii Vitt & Crum, 560; Studies in Cryphaeaceae I. A revision of the genus Cryphaea in North America north of Mexico, 144, II. A review of the monotypic genus Dendropogonella Britt., 521
Marchantia berteroana, 534; palmata, 428, 430; polymorpha, 364, 406, 428 ff., 572
Marchantia polymorpha, Occurrence of androgynous receptacles in, 427
Marquesas Islands, French Polynesia, Mosses of, 174
Marsupella apiculata, 443
Marsupidium flavicans, 534; urvilleanum, 534
Meesea trifaria, 407; uliginosa, 407
Megaceros endivaefolius, 529; fuegiensis, 534
Meiothecium boryanum, 494; negrense, 494; revolubile, 494
Menegazzia terebrata, 209
Merceya ligulata var. bartramii, 193
merochlorophaeic acid, 370
Mesodon onustus, 493
Mesoptychia sahlbergii, 432, 564
metabolic inhibitors, 361
Meteoriopsis patula, 344, 487; recurvifolia, 487; remotifolia, 487, var. robusta, 487; tovariensis, 487
Meteorium auricosta, 486; illecebraria, 487; illecebrum, 487; penicillatum, 487; pseudosinuatum, 487
method for the identification of usnic acid, A specific, 208
methyl hypoprotocetrarate, 79
methyl notate, 79
methyl 2′,-O-methylnotate, 79
methyl 2′,4-di-O-methylhypoprotocetrarate, 79
methyl 4-O-demethylnotate, 79
methyl 4-O-methylhypoprotocetrarate, 79
Metzgeria, Further notes on the production of vegetative thallus structures by female involucres of, and a new species of, 566
Metzgeria litoralis sp. nov. and Apometzgeria from southern South America, 293
Metzgeria acuminata, 566 ff.; **agnewii** sp. nov., 566, 569 ff.; arborescens, 566 ff.; brevialata, 295; decipiens, 534; divaricata, 534; filicina, 566 ff.; frontipilis, 295; fruticulosa, 443; furcata, 566 ff.; grollei, 570; hispidissima, 295; imberbis, 566; liebmanniana, 566 ff.; **litoralis** sp. nov., 293 ff.; macveanii, 293; pinnata, 293, 294; robinsonii, 295; scobina, 570; subscobina, 570; violacea, 534
Mexico, Additional mosses from Costa Rica and, 447; Additions to the moss flora of, 450; and South America, Neosharpiella, a new genus of Musci from high elevations in, 536; Two new lichens (Ramalina) from Baja California, 211
MEYER, MARVIN W., and CARL ANGERMAN: Enzymes in the moss Funaria hygrometrica: phenylalanine ammonia lyase and its control, 361
Micarea lignaria, 66; melaena, 66, 401
Microdus brasiliensis, 478; densus, 478; globicarpus, 478; rostratus, 478
Microphiale pineti, 37
Microthamnium thelistegum, 495; viscidulum, 497; volvatum, 497
Mielichhoferia canescens, 482; graciliseta, 482; gymna, 482
Mittenothamnium diminutivum, 344, 495; elegantulum, 495; langsdorffii, 495; lehmannii, 495; macroblepharum, 176; macrodontium, 495; minusculifolium, 495; nanopolymorphum, 495; reptans, 344, 495; subobscurum, 495
Mniodendron tahiticum, 175
Mniomalia viridis, 482
Mnium spinosum in North America, The occurrence of, 430
Mnium affine, 254 ff., 431, 433; andrewsianum, 432; blyttii, 564; carolinianum, 141; cinclidioides, 562; drummondii, 562; insigne, 431; laevinerve, 205, 206; ligulatum, 482; longirostrum, 482; marginatum, 564; menziesii, 431; orthorrhynchum, 433, 564; riparium, 445, 564; spinosum, 430 ff., 564; spinulosum, 431; stellare, 205, 206, 562; thomsonii, 562; venustum, 431
Moerckia blyttii, 405, 506; hibernica, 315
Mosses from Costa Rica and Mexico, Additional, 447; of North America north of Mexico, A new list, 85; of the Marquesas Islands, French Polynesia, 174; of Utah new to science, 286; Recent literature on, 316, 456, 578
Moss flora of Arizona, Additions to, 439; of Mexico, Additions to, 450; of Texas, and a new species of Isopterygium, 440
Musgos de Venezuela, Un censo de, 473
Mycoblastus affinis, 66, 394; alpinus, 66; sanguinarius, 66

Mycocalicium praecedens, 69
Mycoporum quercus, 40
Mylia anomala, 405
Myurella apiculata, 507; careyana, 563; julacea, 407, 507; siberica, 505, 507, 563, 564; tenerrima, 507
Myuroclada maximowiczii, 446

NAIDU, T. R. B.: Occurrence of androgynous receptacles in Marchantia polymorpha, 428
Nardia geoscyphus, 405; insecta, 443
NASH, THOMAS H. III: Sensitivity of lichens to sulfur dioxide, 333; Two new species of Xanthoparmelia from Arizona, 214; Additions to the lichen flora of Arizona, 545
Neckera complicata, 493; composita, 489; cylindricaulis, 493; denticulata, 487; distans, 493; filiformis, 149, 151; jamesonii, 488; julacea, 497; korthalsiana, 488; moritziana, 497; moritzii, 488; pachygaster, 487; rufescens, 488; squarrosa, 493; suborthosticha, 486; viridis, 487
Neckeropsis disticha, 344, 488; undulata, 344, 488
NELSON, PATRICIA P.: Leptodon smithii (Musci, Neckeraceae), a genus new to North America, 434
Neohypnella chrysophyllopodia, 491; diversifolia, 491
Neosharpiella, a new genus of Musci from high elevations in Mexico and South America, 536
Neosharpiella aztecorum gen. et sp. nov., 536 ff.; **turgida** comb. nov., 537 ff.
Nephroma arcticum, 67, 393; bellum, 547; expallidum, 67, 393
Neuropogon lambii, Rediscovery of, 305
New and interesting macrolichens of British Columbia, 366
Newfoundland, Canada, Buxbaumia aphylla Hedw. in, 187
New list of mosses of North America north of Mexico, 85
New records and comments on lichens of California, 209
News and notes, 224, 331, 472, 593
Nitrogen fixation on granite outcrop pioneer ecosystems, 196
Nomenclatural notes on some members of Parmelia section Melanoparmelia, 306; On the Parmelia xanthina complex, 541
norlobaridone, 307
norobtusatic acid, 77, 78, 81, 83
norstictic acid, 20 ff., 79, 230, 232, 235, 238, 242, 244, 265, 306, 385, 386
North America, Cladonia yunnana new to, 199; Leptodon smithii (Musci, Neckeraceae), a genus new to, 434; Physcia luganensis new to, 421; The occurrence of Mnium spinosum in, 431; The lichen genus Coccotrema in, 260
North America north of Mexico, A new list of mosses of, 85; The genus Lobaria in, 225
North Carolina, Cladonia psoromica, a new lichen species from, 418
Northwest Territories, Canada, Lichens of the Reindeer Preserve, 48
notatic acid, 77, 78, 79, 80, 83
Noteroclada confluens, 534
Notes on some Hawaiian species of the Pottiaceae, 192; on Texas mosses II, 557
Nova Scotia, Scapania massalongoi in—First report from North America, 427
Nowellia curvifolia, 572

obtusatic acid, 77, 78, 81, 83
Occurrence of androgynous receptacles in Marchantia polymorpha, 428
Ocellularia actinota, 79; albogilva, 79; capensis var. feracior, 79, var. leiothallina, 79; gracilis, 80, var. laevigata, 80; granulifera, 80; henatomma, 80; leptopogon, 80; majorina, 80; microporellum, 80; phaeotropa, 80; rhabdospora fo. laevigata, 80; subterebrata, 80; triglyphica, 80
Ochrobryum gardneri, 478
Ochrolechia androgyna, 67, 396; frigida, 67; gemminipara, 396; gibatides, 67; groenlandica, 67; gyalectina, 67, 74; inaequatula, 67; lapuensis, 67; pacifica, 260, 269; parella, 13; pterulina, 67; upsaliensis, 67
Octoblepharum albidum, 175, 344, 478, var. albidum, 478, var. violascens, 478; ampullaceum, 478; cocuiense, 478; cylindricum, 478; erectifolium, 478; longifolium, 175; mittenii, 478; pulvinatum, 447, 449, 478; stramineum, 478; tatei, 478
OHLSSON, KARL E.: New and interesting macrolichens of British Columbia, 366
Oligotrichum hercynicum, 408, 507; parallelum, 507
olivetoric acid, 380, 541
Omphalina ericetorum, 67; hudsonica, 67; luteolilacina, 67
Oncophorus virens, 406; wahlenbergii var. compactus, 446
On the functioning of stomates in Funaria, 423
Oreas martiana, 432
orsellinic acid, 83
Orthocaulis floerkei, 405
Orthodontium fendleri, 482; pellucens, 482
Orthorrhynchium cylindricum, 176
Orthostichella subpachygaster, 488
Orthostichidium excavatum, 486; guyanense, 486
Orthostichopsis auricosta, 486; crinita, 486; tetragona, 486
Orthothecium chryseum, 408; intricatum, 564

Orthotrichum diaphanum, 298; **epapillosum** sp. nov., 437 ff.; holzingeri, 439; jamesianum, 407; laevigatum var. macounii, 445; longifolium, 484; macounii fo. **vermiculare** fo. nov., 291; obtusifolium, 407, 557, 562; pallens, 507; pumilum, 557; pycnophyllum, 557, 558; rivulare, 439; rupestre, 445; sordidum, 562; sprucei, 439; strangulatum, 137, 140, 141; striatum, 445
Orthotrichum from northern California, A new, 437
Oxyrrhynchium clinocarpum, 492; limnobiella, 492

Pachyglossa dissitifolia, 519
Pachyospora verrucosa, 67
Pachyphiale cornea, 36, 37
Palamocladium leskeoides, 492; trichophyllum var. subtile, 492
Pallavicinia lyellii, 571, 572
Paludella squarrosa, 407
Pannaria isidiata, 386; leucosticta, 379, 380; **leucostictoides** sp. nov., 366, 379 ff.; pezizoides, 68, 392
PAOLILLO, DOMINICK J., JR., and DOROTHY L. B. GARNER: A time-course of sporophyte development in Funaria hygrometrica Hedw., 356; On the functioning of stomates in Funaria, 423
PAOLILLO, DOMINICK J., JR., and LEE B. KASS: The germinability of immature spores in Polytrichum, 163
Papillaria aongstroemiana, 176; aongstroemii, 176; crocea, 497; denticulata, 487; deppei, 487; helictophylla, 176; imponderosa, 344, 487; laevifolia, 487; nigrescens, 345, 487, 497, var. illecebra, 487; penicillata, 487; pseudofunalis, 487; seriata, 487; subsquamatula, 487
Paraleucobryum albicans, 478
parietin, 386
Parmelia aberrans, 541, 542; albertana, 397; almquistii, 380, 387; alpicola, 372; **applicata** comb. nov., 307; arseneana, 207; **attica** comb. nov., 306; aurulenta, 547; austerodes, 372; baumgartneri, 306; borreri var. stictica, 382; caperata, 7, 8, 335, 336, 338, 549, var. madagascariacea, 542; ceratophylla var. atrofusca, 372, var. tubulosa, 379; chlorochroa, 68, 74, 84; columnata, 80; conspersa, 33, 38, 196, 197, 198; corallinoides var. pulvinata, 68; cumberlandia, 207; delisei, 306; **dissensa** sp. nov., 214 ff.; domokosii, 80; encrustans, 80, 214; enteromorpha, 374; erecta, 27; exasperatula, 68, 397; flaventior, 397; glomellifera, 308, 309; **huachucensis** sp. nov., 214, 215; hypoleucina, 22, 25, 27; hypopsila, 207; hypotropa, 22, 25, 28; hypotropoides, 27; incurva, 68, 548, 549; infumata, 68, 397; isidiotyla, 308, 309; kurokawae, 215, 547; louisianae, 25; loxodes, 308, 309; madagascariacea, 541, 542, 543; metaphysodes, 378; mexicana, 207, 547; neocongensis, 80, 214; notata, 77, 78, 84; omphalodes, 68; **parahypotropa** sp. nov., 20, 25; perforata, 20 ff.; perlata var. cetrarioides, 370; plittii, 8, 207, 547; praetermissa, 68; **preperforata** sp. nov., 20, 25 ff.; proboscidea var. xanthina, 542; prolixa, 306, 307, var. applicata, 307, var. attica, 306; pseudophysodes, 378; pseudosulcata, 380; psoromifera, 215, 547; revoluta, 380; rigida, 25, 28; rudecta, 335, 336; saubinetii, 382; saxatilis, 9, 68, 380, 382; separata, 68, 74; septentrionalis, 68, 397; sinuosa, 548, 549; sorediosa, 68; sphaerosporella, 207, 210; squarrosa, 382, 387; stictica, 382; stygia, 68; subargentifera, 307; subaurifera, 68, 307, 397, var. subglabra, 307; subdomokosii, 80, 214; **subglabra** comb. nov., 307; subobscura, 379; subolivacea, 398; subrudecta, 398; sulcata, 7, 8, 68, 333, 398; tasmanica, 215, 547; tinctina, 547; tubulosa, 210; verruculifera, 307, 308, 309; weberi, 77, 78, 80, 214; wyomingica, 68; xanthina, 541, 542, 543
Parmelia section Melanoparmelia, Nomenclatural notes on some members of, 306
Parmelia xanthina complex, Chemical and nomenclatural notes on, 541
Parmeliopsis aleurites, 68, 398; ambigua, 68, 210, 398; hyperopta, 68, 398, 547
Pedinophyllopsis abditus, 516, 520
Pellia epiphylla, 503, 504, 572
Peltigera canina (L.) Willd. from sun and shade habitats, photosynthetic pigments in, 543
Peltigera aphthosa, 68, 199, 392; canina, 33, 68, 199, 292, 543, 544, 545; collina, 547; horizontalis, 392; lepidophora, 68; leucophlebia, 68, 392; malacea, 68, 292; membranacea, 69; polydactyla, 69, 392; occidentalis, 383, 387; pulverulenta, 383, 387; praetexta, 392; rufescens, 39, 69, 392; scabrosa, 69, 382, 383, 387; scutata, 387; spuria, 69, 392; venosa, 69, 392
Peltula cylindrica, 196, 197, 198; zahlbruckneri, 547
Pennsylvania, Distributional records of rare or infrequently collected species of mosses, 560
Perforaria minuta, 260, 266
perlatolic acid, 308, 309, 370
Pertusaria carnea, 260, 267, 268; coriacea, 69; curcurbitula, 260, 262, 265, 266; dactylina, 69; glomerata, 69; leioterella, 33; panyrga,

69; pertusa, 38; pocillaria, 267; subobducens, 69; trachydactyla, 69, 74
Phaeocalicium praecedens, 69
Phaeographis eulectra, 210; inusta, 210
Phaeotrema glyphicum, 80; leiostomum, 80; leucocarpum, 80
Phascum cuspidatum, 561
Phenylalanine ammonia lyase and its control, Enzymes in the moss Funaria hygrometrica, 361
phenylalanine ammonia lyase, 361 ff.
Philonotis americana, 565; elongata, 447, 449; filiformis, 483; fontana, 407, 565, var. falcata, 445; glaucescens, 345; gracillima, 483; jardinii, 175; longiseta, 483; macrodictya, 483; moritziana, 483; sphaericarpa, 345, 483; tenella, 483; uncinata, 483, var. glaucescens, 483, var. gracillima, 483; vescoana, 175
Phlyctis agelaea, 36, 37
phosphomonoesterase, 554
photoperiod, temperature, and plant growth hormones, Effect of, on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
Photosynthetic pigments in Peltigera canina (L.) Willd. from sun and shade habitats, 543
Phyllodrepanium falcifolium, 482
Phyllogonium fulgens, 345, 488; goebelii, 488; speciosum, 488; viscosum, 488
Physcia luganensis new to North America, 421
Physcia adscendens, 69, 400, 547; aipolia, 400, var. angustata, 69; albicans, 547; biziana, 547; caesia, 69; ciliata, 400; constipata, 400; dubia, 69; endococcinea, 207; intermedia, 69, 548; luganensis, 421, 422; millegrana, 8, 333, 335, 336, 337, 400; muscigena, 69, 400; nigricans, 422; orbicularis, 400, 421, 422; pusilloides, 421; sciastra, 69; stellaris, 207, 400
Physcomitrium pyriforme, 558; turgidum, 536 ff.
Physconia grisea, 207; muscigena, 69
physodalic acid, 371, 372, 374, 375
physodic acid, 371, 372, 374, 375
Pigafettoa crenulata, 513, 514
Pigafettoa Mass., Austral Hepaticae I, 511
Pilophoron acicularis, 210; nigricaulis, 383
Pilophorus strumaticus, The trichogynes of, 414
Pilopogon gracilis, 478
Pilosium chlorophyllum, 493; flaccisetum, 493
Pilotrichella flexilis, 345, 487; hexasticha, 345; illecebraria, 487; pachygaster, 487; palisotii, 487; pentasticha, 487; rigida, 487; subheterocladia, 488; viridis, 487
Pilotrichidium brunnescens, 345
Pilotrichum affine, 345; amazonum, 489; andersonii, 489; evanescens, 489; fendleri, 489; filiferum, 488; filiforme, 149, var. squarrosulum, 486; heteromalla var. filiforme, 152; hornschuchii, 486; lancifolium, 487; longicaule, 489; nigrescens, 497; nivale, 485; paraphysale, 487; ramosissimum, 489; rufescens, 525; seriatum, 487; turgidum, 487
Pioneer ecosystems, Nitrogen fixation on granite outcrop, 196
Pireella angustifolia, 486; cavifolia, 486, 559; cubensis, 345; cymbifolia, 345, 486; fendleri, 486; filicina, 345; pachyclada, 345
Placopsis cribellans, 263
Placynthium aspratile, 70; nigrum, 70
Plagiochila asplenioides, 405, 572; duricaulis, 534; filipendula, 534; heteromalla, 534; parvidens, 534
Plagiomnium cuspidatum, 407; drummondii, 562; ellipticum, 407
Plagiothecium laetum, 408; schraderi, 493
Plant growth hormones, Effect of, on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
Platismatia glauca, 398
Platydictya jungermannioides, 506
Platygyrium fuscoluteum, 541
Platyhypnidium muelleri, 297; riparioides, 494
Platysma ochrocarpum, 27
Pleurochaete squarrosa, 140
Pleurorthotrichum nukahivense, 175
Pleurozium schreberi, 258, 408
Poecilophyllum procumbens, 477; tortellum, 478
Pogonatum alpiniforme, 496; alpinum var. brevifolium, 565; brachyphyllum, 558; campylocarpon, 496; contortum, 447, 449; flexuosum, 345, 496; laterale, 176; liebmannianum, 496; neesii, 178; oligodus, 496; plurisetum, 496; polycarpum, 496; robustum, 345, 496; semipellucidum, 496; stevensii, 178; subbifarium, 496; tortile, 345, 496; urnigerum, 408
Pohlia annotina, 407, 507, var. loeskei, 447, 449, 507; bulbifera, 507; cruda, 407; cruegeri, 496; longipedicellata, 482; nutans, 190, 407; peracuminata, 447, 449; proligera, 507; rothii, 565; spectabilis, 482
Pollution and drought hypotheses in relation to lichens and bryophytes in urban environments, 1
Polyblastia bryophila, 70; gelatinosa, 70; hyperborea var. macrospora, 70
Polychidium dendriscum, 370, 383, 384
Polytrichadelphus aristatus, 496; flexuosus, 496; nukahivensis, 176; peruvianus, 496; valenciae, 496

Polytrichum antillarum, 496; altisetum var. humilisetum, 473, 496; aristiflorum, 496; brachymitrium, 496; campylocarpum, 496; commune, 163, 164, 165, 166, 408, 496, 554, 555, 558; flexuosum, 496; gracile, 444, var. anomalum, 444; insularum, 345; juniperinum, 163, 164, 166, 168, 190, 345, 408, 473, 496, var. gracilis, 408; ohioense, 168; piliferum, 408; polycarpum, 496; semipellucidum, 496
Polytrichum commune, Electrophoretic study of acid phosphatases in, 554
Polytrichum, The germinability of immature spores in, 163
Porella pinnata, 572; platyphylla, 572
Porothamnium fasciculatum, 488; filiferum, 488
Porotrichodendron robustum, 488; superbum, 488
Porotrichum alleghaniense, 563; flagellare, 488; korthalsianum, 488, 559; longirostre, 488; plicatulum, 488
Potamium casiquiariense, 494; deceptivum, 494; pacimoniense, 494; vulpinum, 494
Pottiaceae, Notes on some Hawaiian species of, 192
Pottia wagneri, 480
Preissia quadrata, 406, 572
Prionodon densus, 486, var. luteovirens, 345; luteovirens, 486; lycopodium, 486; simplex, 486; subgeniculatus, 486
Production of vegetative thallus structures by female involucres of Metzgeria, and a new species of Metzgeria, 566
Protoblastenia monticola, 70; rupestris, 70, 400
protocetraric acid, 375, 380, 549
protolichesteric acid, 299
protolichesterinic acid, 541, 542
Pseudobryum cinclidioides, 562
Pseudocephalozia quadriloba, 534
Pseudocryphaea flagellifera, 345, 486
Pseudocyphellaria anomala, 384, 385; anthraspis, 384, 385; crocata, 210, 385; mougeotiana, 386; rainierensis, 386
Pseudoleskea arizonae, 290
Pseudoleskeella nervosa var. sibirica, 445; williamsii, 445, 565
Pseudopohlia didymodontia, 482
Psoroma hypnorum, 70, 392
psoromic acid, 214, 420, 421
Pterigynandrum cylindricaule, 493; sharpii, 141
Pterobryon densum, 486; fendleri, 486; hornschuchii, 486
Pterogonidium pulchellum, 494
Pterygoneurum rosae, 275
Ptilidium ciliare, 405; pulcherrimum, 405, 572
Ptilium cristacastrensis, 258
Ptychomitrium incurvum, 558
Ptychomnion aciculare, 176
Puiggariella aurifolia, 495
pulvinic acid, 385, 386
pulvinic dilactone, 385, 386
PURSELL, R. A.: Contribución a la brioflora de Venezuela II, 559; Review of E. V. Watson, Mosses, 223; Un censo de los musgos de Venezuela, 473; Distributional records of rare or infrequently collected species of mosses in Pennsylvania, 560
Pycnothelia papillaria, 541
Pylaisiella polyantha, 408, 506
Pyrenopsis phaeococca, 196, 197, 198
Pyrenula leucoplaca, 401; nitida, 40, 42, 401

Quantitative study of Aloina, Aloinella, and Crossidium (Musci), 301

Radula complanata, 572; flavifolia, 534; polyclada, 443
Ramalina almquistii, 70, 74; anceps, 212; bajacalifornica, 212, 213; dasypoga, 212, 213; dendriscoides, 211, 212, 213; druidarum, 80; farinacea, 7; fastigiata, 9, 399; hypoprotocetrarica, 81; intermedia, 70; minuscula, 70, 399; **moranii** sp. nov., 211, 212; obtustata, 70, 74; roesleri, 399; siliquosa, 83, 84; sinensis, 70, 74; thrausta, 399; **wigginsii** sp. nov., 211, 213
Ramalina from Baja California, Mexico, 211
RAMSAY, HELEN P.: Unusual sporocytes in Dicnemoloma pallidum (Hook.) Wijk & Marg., 178
RAO, DHRUVA, and FABIUS LEBLANC: Evaluation of the pollution and drought hypotheses in relation to lichens and bryophytes in urban environments, 1
Rauiella firmula, 491; subcatenulata, 491
RAWAT, M. S., and R. N. CHOPRA: In vitro production of secondary gemmae on the protonema of Bryum klinggraeffii Schimp., 183
Reboulia hemisphaerica, Effect of photoperiod, temperature, and plant growth hormones on initiation of archegoniophore elongation, 501
Reboulia hemisphaerica, 501 ff., 572
Recent literature on hepatics, 452, 574
Recent literature on lichens, 217, 321, 464, 584
Recent literature on mosses, 316, 456, 578
REDFEARN, PAUL L., JR.: Additions to the moss flora of Texas, and a new species of Isopterygium, 440; Review of David W. Shimwell, The description and classification of vegetation, 590
REDHEAD, SCOTT A.: Observations on the rhizoids of Andreaea, 185

Rediscovery of Neuropogon lambii, 305
R[EESE], W. D.: Editorial, 216, 573; Review of Clair A. Brown, Wildflowers of Louisiana and adjoining states, 328; Review of Robert W. Schery, Plants for man, 471; News and notes, 224, 331, 472, 593
Reproductive biology of Riccia crystallina, In vitro studies on, 278
Review of the monotypic genus Dendropogonella Britt., 521
Reviews: K. W. Allison and John Child, The mosses of New Zealand, 329; Clair A. Brown, Wildflowers of Louisiana and adjoining states, 328; Arthur Cronquist, Arthur H. Holmgren, Noel H. Holmgren, and James L. Reveal, Intermountain flora. Vascular plants of the intermountain West, U.S.A., 592; Eilif Dahl and Hildur Krog, Macrolichens of Denmark, Finland, Norway and Sweden, 589; J. H. Dickson, Bryophytes of the Pleistocene. The British record and its chorological implications, 470; A. J. Grout, Moss flora of North America north of Mexico, 470; Yoshiwo Horikawa, Atlas of the Japanese flora: An introduction to plant sociology of East Asia, 471; K. Karczmarz, A monograph of the genus Calliergon (Sull.) Kindb., 327; J. Levitt, Responses of plants to environmental stresses, 328; Geneva Sayre, Cryptogamae Exsiccatae—An annotated bibliography of published exsiccatae of Algae, Lichenes, Hepaticae, and Musci. IV. Bryophyta, 326; Robert W. Schery, Plants for man, 471; David W. Shimwell, The description and classification of vegetation, 590; Edward G. Voss, Michigan flora. Part I. Gymnosperms and monocots, 591; E. V. Watson, Mosses, 223; William A. Weber and Patricia Nelson, Random access key to the genera of Colorado mosses, 222
Rhabdoweisia fugax var. tennerima, 478
Rhacocarpus cuspidatulus, 485; humboldtii, 486; purpurascens, 486
Rhacomitrium aciculare, 562; aquaticum, 444; canescens, 406, var. epilosum, 445, var. ericoides, 406; crispipilum, 480; crispulum, 480; cucullatifolium, 480; heterostichum, 406, var. sudeticum, 406; lanuginosum, 189; ramulosum, 445
Rhacopilopsis trinitensis, 495
Rhacopilum convolutaceum, 176; microphyllum, 176; tomentosum, 345, 485
Rhamphidium fendleri, 480
Rhaphidorrhynchium amoenum, 494; decumbens, 494; decurvifolium, 494; erythropodium, 494; glaziovii, 494, 497; liliputianum, 345; obliquerostratum, 494; sanguisetum, 397, 494; schlimii, 494; subscabrum, 494; trichostegum, 494, 497; xylophilum, 345
Rhaphidostegium richardi, 495
Rhaphidostichum elegans, 494; **pustulatum** sp. nov., 310, 312 ff.
Rhaphidostichum pustulatum, new species of the Sematophyllaceae from Hawaii, 310
Rhizocarpon badioatrum, 70, 548; chioneum, 70; disporum, 70; eupetraeoides, 70; ferax, 70; geographicum, 394, subsp. frigidum, 70, subsp. geographicum, 70; grande, 70, 394, 548; hochstetteri, 70; intermediellum, 548; macrosporum, 207; petraeum, 38; riparium subsp. lindsayanum, 70; superficiale subsp. boreale, 70; tinei subsp. vulgare, 70
Rhizogonium lindigii, 447, 449, 483; spiniforme, 345, 483
Rhizoids of Andreaea, Observations on, 185
Rhizomnium gracile, 407; nudum, 507; pseudopunctatum, 407
Rhizonic acid, 83
Rhodobryum andino-roseum, 482; beyrichianum, 482; domingense, 482; ontariense, 205, 206, 508; pycnopyxis, 482; roseum, 205, 206, 407, 508
Rhynchostegiella compacta, 254 ff., 266, 447
Rhynchostegiopsis flexuosa, 345
Rhynchostegium argute-serratum, 492; beskeanum, 492; campylocarpum, 492; conchophyllum, 492; frondicola, 492; limnobiella, 492; scariosum, 492; sellowii, 492; semiscabrum, 447, 449; serrulatum, 345, 558, var. frondicola, 492; subperspicuum, 493; trachynotum, 493
Rhytidium rugosum, 408
Rhytisma acerinum, 13
Ricasolia ravenelii, 247; tenuis, 244
Riccardia autoica, 534; floribunda, 534; georgiensis, 534; latifrons, 406; palmata, 572; pinguis, 255 ff., 406, 443, 564, 572; prehensilis, 534; tenax, 534; thaxteri, 534
Riccia crystallina, In vitro studies on the reproductive biology of, 278
Riccia crystallina, 278 ff.; frostii, 284; lutescens, 285; robusta, 285
Rigodium gracile, 489; solutum, 489
Rinodina archaea, 58, 71, 399; archaeoides, 71; bischoffii, 71, 399; cinerovirens, 71; lecideoides, 71; lignaria, 71; metaboliza, 71; marysvillensis, 210; mniaraea, 71, 399; mucronulata, 71; nimbosa, 71; pyrina, 399; turfacea, 58, 71, 399
ROBINSON, HAROLD, and C. DELGADILLO M.: Neosharpiella, a new genus of Musci from high elevations in Mexico and South America, 536; and A. J. SHARP and FRANK D. BOWERS: Additional mosses from Costa Rica and Mexico, 447

Roccella montagnei, 33, 38
Roivainenia jacquinotii, 534
ROSENSTRETER, R., and R. P. SHERIDAN: The effect of hydrogen ion concentrations in simulated rain on the moss Tortula ruralis (Hedw.) Sm., 168
RUNDEL, PHILIP W., and PETER A. BOWLER: Two new lichens (Ramalina) from Baja California, Mexico, 211

Saccobasis polita, 405
salazinic acid, 21, 29, 211, 380, 382, 549
Sarcogyne clavus, 196 ff.; novomexicana, 548; pruinosa, 71; regularis, 71, 396
scabrosin, 383
Scapania apiculata, 427; curta, 405, 427; degenii, 571, 572; glaucocephala, 427; irrigua, 405; massalongoi, 405; nemorosa, 572; paludosa, 564; perssonii, 572, 573; uliginosa, 564
Scapania massalongoi in Nova Scotia—First report from North America, 427
Scapania perssonii Schust., A note on, 572
Schistocheila berteroana, 534; laminigera, 534; reflexa, 534
Schlotheimia glauca, 485; grandiareolata, 485; jamesonii, 485; lasiomitria, 485; merkelii, 485; pungentissima, 485; rugifolia, 485; torquata, 345, 485; trichomitria, 485
Schraderella pungens, 494
SCHROEDER, GARY J., DOYLE E. ANDEREGG, and NANCY E. SCHROEDER: Cladonia yunnana new to North America, 199; Further additions to the lichen flora of Idaho, 207
SCHUSTER, R. M.: A note on Scapania perssonii Schust., 572; and JOHN J. ENGEL: Austral Hepaticae I. Pigafettoa Mass., 511; II. Evansianthus, a new genus of Geocalycaceae, 516
Scopelophila cataractae, 193; **infericola** sp. nov., 192, 194 ff.; ligulata, 193
Scorpidium scorpioides, 408, var. venezuelense, 492; turgescens, 408
SCOTTER, GEORGE W.: Lichens of the Reindeer Preserve, Northwest Territories, Canada, 48
scrobiculin, 230, 232, 238, 251
secondary gemmae, In vitro production on the protonema of Bryum klinggraeffii, 183
Second report for Drummondia pursellii Vitt & Crum, 560
Seligeria globicarpa, 478; rostrata, 478
Sematophyllum adnatum, 558; arrectum, 495; caespitosum, 345; chrysostegum, 494; constrictum, 494; cuspidiferum, 494; erythropodium, 494; flaccidifolium, 494; galipense, 345, 447, 449, 494; glaziovii, 497; inundatum, 495; kegelianum, 494; lamprophyllum, 345; loxense var. lithophilum, 494; sanguisetum, 497; sericifolium, 345; subsimplex, 495; tequendamense, 495; trichostegum, 497
Semibarbula orientalis, 203, 205
Sensitivity of lichens to sulfur dioxide, 333
SHARP, A. J.: Review of Yoshiwo Horikawa, Atlas of the Japanese flora; An introduction to plant sociology of East Asia, 471; and FRANK D. BOWERS and HAROLD ROBINSON: Additional mosses from Costa Rica and Mexico, 447
SHERIDAN, R. P., and R. ROSENSTRETER: The effect of hydrogen ion concentrations in simulated rain on the moss Tortula ruralis (Hedw.) Sm., 168
Siphula simplex, 210
SNYDER, J. M., and L. H. WULLSTEIN: Nitrogen fixation on granite outcrop pioneer ecosystems, 196
Solenostoma crenulatum, 572; pumilum, 572
Solmsiella biseriata, 484
Solorina bispora, 71; crocea, 71, 393; saccata, 71, 393; spongiosa, 71
Solorinella asteriscus, 388, 393
SOOD, SNEH, and R. N. CHOPRA: In vitro studies on the reproductive biology of Riccia crystallina, 278
South America, Metzgeria litoralis sp. nov. and Apometzgeria from, 293; Neosharpiella, a new genus of Musci from high elevations in, 536; The Raymond E. Hatcher collection of Hepaticae and Anthocerotae from southern, 528
Southern Appalachian Mountains, Two lichens (Parmeliae) new to, 548
Sphaerophorus globosus, 71, 207
Sphagnum acutifolium var. meridense, 474; alegrense, 474; angustifolium, 509; balticum, 508; capillaceum, 255, 256, 257, 258, 259, 406, 508, var. tenellum, 474, 508; centrale, 508; compactum, 508; coryphaeum, 474; cuspidatum, 509; erythrocalyx, 474; flavicaule, 474; fuscum, 406, 508; girgensohnii, 508; gracilescens, 474; limbatum, 474; lindbergii, 508; magellanicum, 406, 474, 508; majus, 508; medium, 474; meridense, 474; molle, 443; obtusum, 508, 509; oxyphyllum, 474; palustre, 474; portoricense, 474; recurvum, 474, 508, 509, var. tenue, 406, 508, 509; riparium, 508; rubellum, 474; russowii, 508; sanguinale, 474; squarrosum, 508; strictum, 474; subsecundum, 474, 508, var. rufescens, 558; tenerum, 508; teres, 508; warnstorfii, 390, 406, 508; wulfianum, 508
Splachnobryum obtusum, 296, 297, 481
Splachnum ovatum, 407; vasculosum, 351, 352; wormskioldii, 347, 349
Sporastatia testudinea, 548

Sporophyte development in Funaria hygrometrica Hedw., A time-course of, 356
Squamaria lentigera, 396
squamatic acid, 200, 420, 421
Squamidium brasiliense, 488; filiferum, 488; leucotrichum, 345, 488; nigricans, 345, 487, 488, 559, var. pendula, 473, 488; subheterocladium, 488
Staurothele clopima, 71; fuscocuprea, 71; perradiata, 71
STEERE, WILLIAM CAMPBELL: Observations on the genus Aplodon (Musci: Splachnaceae), 347; The occurrence of Mnium spinosum in North America, 430; Review of J. H. Dickson, Bryophytes of the Pleistocene, 470; Review of A. J. Grout, Moss Flora of North America north of Mexico, 470; and HOWARD A. CRUM and LEWIS E. ANDERSON: A new list of mosses of North America north of Mexico, 85
Stenodictyon pallidum, 345
stenosporic acid, 306, 308
Stereocaulon alpinum, 71, 394, var. erectum, 72; condensatum, 72; evolutum, 210; glareosum, 72; intermedium, 210; myriocarpum, 210; paschale, 72; rivulorum, 72; sterile, 210; tomentosum, 72, 210, 394
Stereophyllum cultelliforme, 345, 493; leucostegum, 345, 493; obtusum, 493; papilliferum, 493; radiculosum, 345, 451, 493
Steyermarkiella anomalodictya, 478
Sticta fuliginosa, 545, 548; hallii, 238; linita var. ochroleuca, 241; mougeotiana, 386; oregana, 241; ravenelii, 247; wrightii, 251
stictic acid, 20 ff., 230, 232, 233, 235, 238, 244, 265, 268, 383, 386
Stictina hallii, 238
stomates in Funaria, On the functioning of, 423
Streptopogon calymperes, 480; cavifolius, 480; erythrodontus, 447, 449; lindigii, 480
STRINGER, PAUL W., and MURIEL H. L. STRINGER: Studies on the bryophytes of southern Manitoba. VII. Distribution of terrestrial bryophytes in a black spruce bog in Bird's Hill Provincial Park, 252
Studies in Cryphaeaceae I. A revision of the genus Cryphaea in North America north of Mexico, 144; II. A review of the monotypic genus Dendropogonella Britt., 521
sulfur dioxide, Sensitivity of lichens to, 333
Symphyogyna hochstetteri, 534
Syrrhopodon banksii, 175; bernoullii, 479; circinatus, 479; compactulus, 479; cryptocarpus, 497; cylindrothecius, 479; epapillosus, 479; fimbriatus, 479; flexiareolatus, 479; floridanus, 345; gaudichaudii, 345, 479; hawaiicus, 297; hioramii, 345; incompletus, 345; kilaueae, 297; leprieurii, 479; lycopodioides, 345, 479; macroprolifer, 479; papillosus, 479; parasiticus, 479; pomiformis, 497; prolifer, 345, 479; rigidus, 479; rubiginosus, 479; rupestris, 479; sordidus, 479; steyermarkii, 479; tenuifolius, 346, 479; venezuelanus, 479; xanthophyllus, 479

Tasmania, Heterodea muelleri in, 410
Taxicaulis andinosubulatus, 492
Taxithelium falcifolium, 176; planum, 346, 495; vernieri, 176
Tayloria lingulata, 406; moritziana, 481; sandwicensis, 298; scabriseta, 481; serrata, 565
Telaranea plumulosa, 534
Teloschistes arcticus, 72, 74, 75; contortuplicatus, 72, 74
Temnoma chilense, 529; pilosum, 534; quadripartitum, 534
temperature, Effect of, on initiation of archegoniophore elongation in the thalloid liverwort Reboulia hemisphaerica, 501
tenuorin, 230, 232, 233, 241, 245, 383, 386
Tetraphis pellucida, 565
Tetraplodon angustatus, 407; mnioides, 407, 481
Texas, Additions to the moss flora of, 440; Notes on mosses II, 557
Thamnium subseriatum, 488
Thamnobryum alleghaniense, 563
Thamnolia vermicularis, 72, var. subuliformis, 72
thelephoric acid, 230, 235, 236, 237, 379
Thelia asprella, 558; hirtella, 558; lescurii, 558
Thelidium absconditum, 72; microsporum, 72; minimum, 72; velutinum, 72
Thelocarpon sphaerosporum, 56
Thelotrema leucohymenium, 81; myrioporoides, 81; occlusum, 81
THOMSON, JOHN W.: Review of Eilif Dahl and Hildur Krog, Macrolichens of Denmark, Finland, Norway and Sweden, 489
Thrombium epigaeum, 401
Thuidiopsis furfurosa, 447, 449
Thuidium acuminatum, 492; allenii, 558; antillarum, 492; delicatulum, 492, var. radicans, 255 ff.; frontinoae, 491; gratum, 491; involvens, 346, 491; minutulum, 491; muriculatum, 491; pauperum, 491; pseudoprotensum, 492; pusillum, 492; recognitum, 140, 254 ff., 407, var. delicatulum, 492; scabrosulum, 492; tamariscinum, 492
Thyridium obtusifolium, 175
Thysanomitrium arctocarpum, 476; caracasanum, 476; luteum, 476; scabrisetum, 476
TIBBETTS, DONALD L., and KATHLEEN H. CHANCE: A specific method for the identification of usnic acid, 208
Time-course of sporophyte development in Funaria hygrometrica Hedw., 356

Timmia austriaca, 407; bavarica, 447, 449; megapolitana var. bavarica, 407
Tomenthypnum nitens, 255 ff., 350, 408
Toninia alutacea, 72; caeruleonigricans, 72, 394, 548; candida, 72, 394; cumulata, 72; lobulata, 72, 394; ruginosa, 394; tristis, 72
Tortella flavovirens, 140; fragilis, 292, 406; humilis, 140, 346, 480, 558; mollissima, 346, 480; tortuosa, 406
Tortula agraria, 346; andicola, 480; bifrons, 273; cucullifera, 276; desertorum, 183, 185; elongata, 480; fragilis, 480; mniifolia, 480; mucronifolia, 406; pagorum, 558; papillosa, 561; rhizophylla, 195; rigida var. pilifera, 273; ruralis, 168 ff., 185, 406; stanfordensis, 183, 185; vectensis, 194, 195
Trematodon fendleri, 475
Tremotylium angolense, 36
Triandrophyllum trifidum, 534
Trichocolea elegans, 534
Trichogynes of Pilophorus strumaticus, 414
Trichosteleum ambiguum, 495; arrectum, 495; cyparissoides, 494; fluviale, 447, 449, 495; inundatum, 495; papillosum, 495; sentosum, 495
Trichostomopsis australasiae, 480
Trichostomum duidense, 480; fendleri, 480; involutum, 346; jamaicense, 480; juniperinum, 448; linealifolium, 480, 497; lineatum, 496; oblongifolium, 195; tovarense, 480; wagneri, 480
Tristichium mirabile, 475
Tritomaria exsecta, 427, 571; exsectiformis, 405, 572; quinquedentata, 405; scitula, 405
Tromera difformis, 34, 37; resinae, 34, 37
TUCKER, SHIRLEY C.: New records and comments on lichens of California, 209
Two bryophytes new to Virginia, 315
Two lichens (Parmeliae) new to the Southern Appalachian Mountains, 548
Two new lichens (Ramalina) from Baja California, Mexico, 211
Two new species of Xanthoparmelia from Arizona, 214

Ulota bruchii, 445; curvifolia, 505, 509
Umbilicaria arctica, 72; cylindrica, 395; hyperborea, 72, 395; papulosa, 552; pennsylvanica, 551 ff.; pertusa, 550 ff.; proboscidea, 72; pustulata, 551 ff.; torrefacta, 72
Umbilicaria pertusa Rass. and some related lichen species, 550
Urban environments, Evaluation of the pollution and drought hypotheses in relation to lichens and bryophytes, 1
Usnea cavernosa, 398; ceratina, 398, 412, 413; compacta, 398; dasypoga, 398; fragilescens, 9; glabrata, 72, 398; glabrescens, 398; hirta, 398; lapponica, 72; scabrata subsp. nylanderiana, 399; subfloridana, 8, 399; substerilis, 399
usnic acid, A specific method for the identification of, 208
usnic acid, 80, 81, 208 ff., 213, 214, 232, 233, 238, 242, 370, 420, 542
Utah mosses new to science, 286

VÄNSKÄ, HEINO, TEUVO AHTI, and GEORGE W. SCOTTER: Lichens of the Reindeer Preserve, Northwest Territories, Canada, 48
Varicellaria kemensis, 73, 395; microsticta, 73; rhodocarpa, 73
Venezuela, Un censo de los musgos de, 473; Contribución a la brioflora de, II, 559
Verrucaria acrotella, 73; aethiobola, 73; cataleptoides, 73; confluens, 40, 42; deversa, 73; devergens, 73; difformis, 40; muralis, 73; scotina, 73; sphinctrina, 73; umbrinula, 73; viridula, 548
Vesicularia amphibola, 493; inflectens, 176; subdenticulata, 494; vesicularis, 346, 494
Vestergrenopsis isidiata, 386
Virginia, Two bryophytes new to, 315
VITT, DALE H.: Distributional studies on bryophytes of Alberta, 505; Species collected in Alberta on the first 1971 foray of the American Bryological and Lichenological Society. Part II. Bryophytes, 403; Review of K. W. Allison and John Child, The mosses of New Zealand, 329; Review of William A. Weber and Patricia Nelson, Random access key to the genera of Colorado mosses, 222
Voitia hyperborea, 432; nivalis, 432
VOTH, PAUL D.: Review of Edward G. Voss, Michigan flora. Part I, 591

Webera papillosa, 482
WEBER, WILLIAM A.: Rediscovery of Neuropogon lambii, 305
Weissia controversa, 131 ff., 178, 346, 444, 451, 480, 558, 561, 565, var. edentula, 143, var. longiseta, 140; ligulifolia, 292; melanostoma, 480; **perligulata** sp. nov., 291; **sharpii** sp. nov., 131 ff.; venezuelensis, 480; viridula, 561
Weissia sharpii, a new species from North America, with cytological and ecological correlations, 131
West Himalayan mosses, Cytological studies on some—III, 202
WEST, STEPHEN D.: Buxbaumia aphylla in Alaska, 565
WHITTIER, HENRY O.: Mosses of the Marquesas Islands, French Polynesia, 174
WILLIAMS, HARRY: Scapania massalongoi in Nova Scotia—First report from North America, 427

Wisconsin, Hepatics from southwestern, 571
WULLSTEIN, L. H., and J. M. SNYDER: Nitrogen fixation on granite outcrop pioneer ecosystems, 196

Xanthoparmelia from Arizona, Two new species of, 214
Xanthoria candelaria, 73; elegans, 73, 400; fallax, 8, 400; hasseana, 386; parietina, 8, 34, 36; polycarpa, 386, 400; sorediata, 73, 548
Xenocephalozia navicularis, 534
Xylographa abietina, 73, 400; vitiligo, 73

Zahlbrucknerella calcarea, 73
zeorin, 383
Zygodon apiculatus, 141; fendleri, 485; gymnus, 485; linguiformis, 485; moritzianus, 485; obtusifolius, 485; pichinchensis, 485; pilosulus, 485; pungens, 485; reinwardtii, 485; subdenticulatus, 485; tetragonostomus, 298

'T was summer, and the spot a cool retreat—
Where curious eyes came not, not footstep rude
Disturbed the lovers' chosen solitude:
Beneath an oak there was a mossy seat,
Where we reclined, while birds above us wooed
Their mates in songs voluptuously sweet.

ELIZABETH CLEMENTINE KINNEY
A Dream

Here from the brow of the hill I look,
Through a lattice of boughs and leaves,
On the old gray mill with its gambrel roof,
And the moss on its rotting eaves.
I hear the clatter that jars its walls,
And the rushing water's sound,
And I see the black floats rise and fall
As the wheel goes slowly round.

THOMAS DUNN ENGLISH
The Old Mill

Then came green meadows, broad and bright,
Where dandelions, with wealth untold,
Gleamed on the young and eager sight
Like stars of gold;
And on the very meadows edge,
Beneath the ragged blackberry hedge,
Mid mosses golden, gray and green,
The fresh young buttercups were seen,
And small spring-beauties, sent to be
The heralds of anemone:
All just as when I earliest heard
The cardinal bird.

WILLIAM DAVIS GALLAGHER
The Cardinal Bird

In vain, in vain; we meet no more,
Nor dream what fates befall;
And long upon the stranger's shore
My voice on thee may call,
When years have clothed the line in moss
That tells thy name and days,
And withered, on thy simple cross,
The wreaths of Père-la-Chaise!

OLIVER WENDELL HOLMES
La Grisette

The woods are greening overhead,
And flowers adorn each mossy bed:
The waters babble as they run—
One thing is lacking, only one:
If Mary were but here today,
I would believe your charming lay,
"Witchery—witchery—witchery!"

HENRY VAN DYKE
The Maryland Yellow-Throat

Volume 76, Number 3 (pages 333–472), was issued 27 September 1973

THE BRYOLOGIST

Quarterly Journal of the American Bryological and Lichenological Society

Editor
WILLIAM D. REESE

Volume 76
1973

Table of Contents

AHTI, T., G. W. SCOTTER & H. VÄNSKÄ. Lichens of the Reindeer Preserve, Northwest Territories, Canada 48

ALMEDA, F., JR. & J. P. DEY. Chemical and nomenclatural notes on the *Parmelia xanthina* complex 541

ANDEREGG, D. E., G. J. SCHROEDER & N. E. SCHROEDER. Further additions to the lichen flora of Idaho 207

ANDERSON, L. E. & B. E. LEMMON. *Weissia sharpii*, a new species from North America, with cytological and ecological correlations 131

BIRD, C. D. Species collected in Alberta on the first 1971 foray of the American Bryological and Lichenological Society. Part I. Introduction and lichens 388

BIZOT, M. Contribution a la flore bryologique de Cuba (2° article) 340

BLACKMAN, A. J., G. C. BRATT & J. A. CASHIN. Distribution and chemistry of *Heterodea muelleri* (Hampe) Nyl. 410

BOWERS, F. D., A. J. SHARP & H. ROBINSON. Additional mosses from Costa Rica and Mexico 447

BOWLER, P. A. & P. W. RUNDEL. Two new lichens (*Ramalina*) from Baja California, Mexico 211

BRASSARD, G. R. Review 327

BREIL, D. A. Two bryophytes new to Virginia 315

BRODO, I. M. The lichen genus *Coccotrema* in North America 260

CHANCE, K. H. & D. L. TIBBETTS. A specific method for the identification of usnic acid 208

CHOPRA, R. N. & S. SOOD. *In vitro* studies on the reproductive biology of *Riccia crystallina* 278

——— & M. S. RAWAT. *In vitro* production of secondary gemmae on the protonema of *Bryum klinggraeffii* Schimp. 183

CROSBY, M. R. Recent literature on mosses—94–96 ..... 316, 456, 578

CRUM, H. Review 590

———, W. C. STEERE & L. E. ANDERSON. A new list of mosses of North America north of Mexico 85

CULBERSON, C. F. & M. E. HALE, JR. 4-O-Demethylnotatic acid, a new depsidone in some lichens producing hypoprotocetraric acid 77

CULBERSON, W. L. The *Parmelia perforata* group: Niche characteristics of chemical races, speciation by parallel evolution, and a new taxonomy 20

———. Recent literature on lichens—83–86 ..... 217, 321, 464, 584

DELGADILLO M., C. A new species, nomenclatural changes, and generic limits in *Aloina*, *Aloinella*, and *Crossidium* (Musci) 271

———. A quantitative study of *Aloina*, *Aloinella*, and *Crossidium* (Musci) 301

DEY, J. P. *Cladonia psoromica*, a new lichen species from western North Carolina 418

———. Two lichens (*Parmeliae*) new to the Southern Appalachian Mountains 548

ENGEL, J. J. The Raymond E. Hatcher collection of Hepaticae and Anthocerotae from southern South America, with new taxa and notes on range extensions 528

——— & Y. KUWAHARA. *Metzgeria litoralis* sp. nov. and *Apometzgeria* from southern South America 293

——— & R. M. SCHUSTER. Austral Hepaticae I. *Pigafettoa* Mass. 511

ESSLINGER, T. L. Nomenclatural notes on some members of *Parmelia* section *Melanoparmelia* 306

———. *Physcia luganensis* new to North America 421
FLORSCHÜTZ, P. A. Review 326
FLOWERS, S. Mosses of Utah new to science 286
FULFORD, M. Recent literature on hepatics—65–66 ..... 452, 574
GARNER, D. & D. J. PAOLILLO, JR. A time-course of sporophyte development in *Funaria hygrometrica* Hedw. 356
——— & ———. On the functioning of stomates in *Funaria* 423
GUERKE, W. R. Epiphyllous Hepaticae in Louisiana 201
GUNDERSON, G. O. Hepatics from southwestern Wisconsin 571
HAMPTON, R. E. Photosynthetic pigments in *Peltigera canina* (L.) Willd. from sun and shade habitats 543
HANCOCK, J. A. & G. R. BRASSARD. *Buxbaumia aphylla* Hedw. in southeastern Newfoundland, Canada 187
HERMANN, F. J. Additions to the bryophyte flora of Alaska 442
———. Additions to the bryophyte flora of Mt. McKinley National Park, Alaska 563
HOE, W. J. Notes on some Hawaiian species of the Pottiaceae 192
———. Additional new and noteworthy records for Hawaiian mosses—2 296
———. *Glossadelphus abortivapicus* and *Rhaphidostichum pustulatum*, new species of the Sematophyllaceae from Hawaii 310
JAHNS, H. M. The trichogynes of *Pilophorus strumaticus* 414
JOHNSEN, A. B. Additions to the moss flora of Arizona. II 439
JORDAN, W. P. The genus *Lobaria* in North America north of Mexico 225
KAHLEM, G. & C. HÉBANT. Electrophoretic study of acid phosphatases in *Polytrichum commune* 554
KERSHAW, K. A. Review 328
KOEVENIG, J. L. Effect of photoperiod, temperature, and plant growth hormones on initiation of archegoniophore elongation in the thalloid liverwort *Reboulia hemisphaerica* 501
KRAL, R. Review 592
KROG, H. *Cetraria inermis* (Nyl.) Krog, a new lichen species in the Amphi-Beringian flora element 299
———. On *Umbilicaria pertusa* Rass. and some related lichen species 550
KUMAR, S. S. Cytological studies on some west Himalayan mosses—III 202
KUWAHARA, Y. Further notes on the production of vegetative thallus structures by female involucres of *Metzgeria*, and a new species of *Metzgeria* 566
LAWTON, E. & F. J. HERMANN. A new *Orthotrichum* from northern California 437
LEBLANC, F. & D. N. RAO. Evaluation of the pollution and drought hypotheses in relation to lichens and bryophytes in urban environments 1
LETROUIT-GALINOU, M.-A. Les Asques des Lichens et la type archaeascé 30
MAGILL, R. E. Notes on Texas mosses II 557
MANUEL, M. G. Studies in Cryphaeaceae I. A revision of the genus *Cryphaea* in North America north of Mexico 144
———. Additions to the moss flora of Mexico—II 450
———. Studies in Cryphaeaceae II. A review of the monotypic genus *Dendropogonella* Britt. 521
———. A second report for *Drummondia pursellii* Vitt & Crum 560
MEYER, M. W. & C. ANGERMAN. Enzymes in the moss *Funaria hygrometrica*: Phenylalanine ammonia lyase and its control 361
NAIDU, T. R. B. Occurrence of androgynous receptacles in *Marchantia polymorpha* 428

NASH, T. H. III. Two new species of *Xanthoparmelia* from Arizona 214
———. Sensitivity of lichens to sulfur dioxide 333
———. Additions to the lichen flora of Arizona I 545
NELSON, P. P. *Leptodon smithii* (Musci, Neckeraceae), a genus new to North America 434
OHLSSON, K. E. New and interesting macrolichens of British Columbia 366
PAOLILLO, D. J., JR. & L. B. KASS. The germinability of immature spores of *Polytrichum* 163
PURSELL, R. A. Review 223
———. Un censo de los musgos de Venezuela 473
———. Contribución a la brioflora de Venezuela II 559
———. Distributional records of rare or infrequently collected species of mosses in Pennsylvania 560
RAMSAY, H. P. Unusual sporocytes in *Dicnemoloma pallidum* (Hook.) Wijk & Marg. 178
REDFEARN, P. L., JR. Additions to the moss flora of Texas, and a new species of *Isopterygium* 440
———. Review 590
REDHEAD, S. A. Observations on the rhizoids of *Andreaea* 185
REESE, W. D. Editorials 216, 573
———. Reviews 328, 471
ROBINSON, H. & C. DELGADILLO M. *Neosharpiella*, a new genus of Musci from high elevations in Mexico and South America 536
SCHROEDER, G. J., D. E. ANDEREGG & N. E. SCHROEDER. *Cladonia yunnana* new to North America 199
SCHUSTER, R. M. A note on *Scapania personnii* Schust. 572
——— & J. J. ENGEL. Austral Hepaticae II. *Evansianthus*, a new genus of Geocalycaceae 516
SHARP, A. J. Review 471
SHERIDAN, R. P. & R. ROSENSTRETER. The effect of hydrogen ion concentrations in simulated rain on the moss *Tortula ruralis* (Hedw.) Sm. 168
SNYDER, J. M. & L. H. WULLSTEIN. Nitrogen fixation on granite outcrop pioneer ecosystems 196
STEERE, W. C. Observations on the genus *Aplodon* (Musci: Splachnaceae) 347
———. The occurrence of *Mnium spinosum* in North America 430
———. Reviews 470
STRINGER, P. W. & M. H. L. STRINGER. Studies on the bryophytes of southern Manitoba VII. Distribution of terrestrial bryophytes in a black spruce bog in Bird's Hill Provincial Park 252
THOMSON, J. W. Review 589
TUCKER, S. C. New records and comments on lichens of California 209
VITT, D. H. Reviews 222, 329
———. Species collected in Alberta on the first 1971 foray of the American Bryological and Lichenological Society. Part II. Bryophytes 403
———. Distributional studies on bryophytes of Alberta 505
VOTH, P. D. Review 591
WEBER, W. A. Rediscovery of *Neuropogon lambii* 305
WEST, S. D. *Buxbaumia aphylla* in Alaska 565
WHITTIER, H. O. Mosses of the Marquesas Islands, French Polynesia 174
WILLIAMS, H. *Scapania massalongoi* in Nova Scotia—first report from North America 427